

Hamilton Geometry

-

The geometry of spacetime from (modified) dispersion relations

Christian Pfeifer
Institute for Theoretical Physics,
Leibniz Universität Hannover

Joint work with L. Barcaroli, L. Brunkhorst, G. Gubitosi, N. Loret

GR21
New York, Juli 2016



Leibniz
Universität
Hannover

1. Dispersion Relations and the Geometry of Spacetime
2. Hamiltonian Phase Space Geometry
3. Conclusion and Outlook

1. Dispersion Relations and the Geometry of Spacetime

2. Hamiltonian Phase Space Geometry

3. Conclusion and Outlook

The dispersion relation and the geometry of spacetime in GR

The dispersion relation of a free point particle in general relativity

$$E^2 - p_\alpha p_\beta \delta^{\alpha\beta} = E^2 - \vec{p}^2 = m^2$$

- m is the invariant mass parameter
- $E = g(\dot{\gamma}, p)$ is the energy of to the particle
- $p_\alpha = g(e_\alpha, p)$ is the spatial momentum of to the particle

an observer on worldline γ associates to the particle with 4-momentum p

The dispersion relation and the geometry of spacetime in GR

The dispersion relation of a free point particle in general relativity

$$E^2 - p_\alpha p_\beta \delta^{\alpha\beta} = E^2 - \vec{p}^2 = m^2$$

- m is the invariant mass parameter
- $E = g(\dot{\gamma}, p)$ is the energy of to the particle
- $p_\alpha = g(e_\alpha, p)$ is the spatial momentum of to the particle

an observer on worldline γ associates to the particle with 4-momentum p

Covariant: The dispersion relation is a level set of a Hamilton function

$$H(x, p) = g^{ab}(x) p_a p_b = m^2$$

The dispersion relation and the geometry of spacetime in GR

The dispersion relation of a free point particle in general relativity

$$E^2 - p_\alpha p_\beta \delta^{\alpha\beta} = E^2 - \vec{p}^2 = m^2$$

- m is the invariant mass parameter
- $E = g(\dot{\gamma}, p)$ is the energy of to the particle
- $p_\alpha = g(e_\alpha, p)$ is the spatial momentum of to the particle

an observer on worldline γ associates to the particle with 4-momentum p

Covariant: The dispersion relation is a level set of a Hamilton function

$$H(x, p) = g^{ab}(x) p_a p_b = m^2$$

The covariant dispersion relation demonstrates

- its intimate relation to the geometry of spacetime, i.e. the metric g
- the geometry of spacetime is derived from second derivatives of H w.r.t. the momenta p of particles
- particle worldlines are determined by Hamilton's equations of motion

The dispersion relation and the geometry of spacetime in GR

Planck scale modified dispersion relation of a free point particle

Doubly Special Relativity [Amelino-Camelia 2008], Relative locality [Amelino-Camelia, Freidel, Kowalski Gilkman, Smolin 2011]

$$E^2 - \vec{p}^2 = m^2 \quad \rightarrow \quad E^2 - \vec{p}^2 + \ell f(E, \vec{p}) + \dots = m^2$$

- What is the underlying spacetime geometry?
- Higher orders in E and \vec{p} cannot yield metric spacetime geometry
- Relation between the particles 4-momentum p and E, \vec{p} ?

The dispersion relation and the geometry of spacetime in GR

Planck scale modified dispersion relation of a free point particle

Doubly Special Relativity [Amelino-Camelia 2008], Relative locality [Amelino-Camelia, Freidel, Kowalski Gilkman, Smolin 2011]

$$E^2 - \vec{p}^2 = m^2 \quad \rightarrow \quad E^2 - \vec{p}^2 + \ell f(E, \vec{p}) + \dots = m^2$$

- What is the underlying spacetime geometry?
- Higher orders in E and \vec{p} cannot yield metric spacetime geometry
- Relation between the particles 4-momentum p and E, \vec{p} ?

Idea: Start from covariant dispersion relation!

Covariant: The dispersion relation is a level set of a Hamilton function

$$H(x, p) = g^{ab}(x)p_ap_b + \ell G^{abc}(x)p_ap_bp_c + \ell^2 H^{abcd}(x)p_ap_bp_cp_d + \dots = m^2$$

The dispersion relation and the geometry of spacetime in GR

Planck scale modified dispersion relation of a free point particle

Doubly Special Relativity [Amelino-Camelia 2008], Relative locality [Amelino-Camelia, Freidel, Kowalski Gilkman, Smolin 2011]

$$E^2 - \vec{p}^2 = m^2 \quad \rightarrow \quad E^2 - \vec{p}^2 + \ell f(E, \vec{p}) + \dots = m^2$$

- What is the underlying spacetime geometry?
- Higher orders in E and \vec{p} cannot yield metric spacetime geometry
- Relation between the particles 4-momentum p and E, \vec{p} ?

Idea: Start from covariant dispersion relation!

Covariant: The dispersion relation is a level set of a Hamilton function

$$H(x, p) = g^{ab}(x)p_ap_b + \ell G^{abc}(x)p_ap_bp_c + \ell^2 H^{abcd}(x)p_ap_bp_cp_d + \dots = m^2$$

The covariant dispersion relation encodes

- the geometry of spacetime, in the tensors g, G, H, \dots
- the geometry of spacetime is derived from second (and higher) derivatives of H w.r.t. the momenta p of particles
- particle worldlines are determined by Hamiltons equations of motion

1. Dispersion Relations and the Geometry of Spacetime

2. Hamiltonian Phase Space Geometry

3. Conclusion and Outlook


Hamilton Geometry

Hamilton phase space geometry

- fundamental tuple (T^*M , $H(x,p)$)
cotangent bundle with
Hamiltonian

Metric spacetime geometry:

- fundamental tuple (M , $g(x)$)
spacetime with metric



Hamilton Geometry

Hamilton phase space geometry

- fundamental tuple (T^*M , $H(x,p)$)
cotangent bundle with Hamiltonian
- the Cartan non-linear connection
 N determines the geometry of
phase space

$$N_{ab} = \frac{1}{4} \left(\{g_{ab}^H, H\} + g_{ai}^H \partial_b \bar{\partial}^i H + g_{bi}^H \partial_a \bar{\partial}^i H \right)$$

$$g^{Hab} = \frac{1}{2} \frac{\partial}{\partial p_a} \frac{\partial}{\partial p_b} H = \frac{1}{2} \bar{\partial}^a \bar{\partial}^b H$$

unique, symmetric, torsion-free


Metric spacetime geometry:

- fundamental tuple (M , $g(x)$)
spacetime with metric
- the lifted Levi-Civita connection
 $p\Gamma$ determines the geometry of
phase space

$$p_a \Gamma^a_{bc} = \frac{1}{2} p_a g^{aq} (\partial_b g_{cq} + \partial_c g_{bq} - \partial_q g_{bc})$$

$$g^{Hab} = g^{ab}$$

unique, symmetric, torsion-free



Hamilton Geometry

Hamilton phase space geometry

- fundamental tuple (T^*M , $H(x,p)$)
cotangent bundle with Hamiltonian
- the Cartan non-linear connection N
determines the geometry of phase space
- the spacetime connection and its
curvature determine the
geometry of spacetime

$$\Gamma^{\delta a}_{bc} = \frac{1}{2} g^{Haq} (\delta_b g_{cq}^H + \delta_c g_{bq}^H - \delta_q^H g_{bc}^H)$$

$$\delta_a = \partial_a - N_{ab}(x, p) \bar{\partial}^b$$

$$R^a_{bcd}(x, p) = 2\delta_{[c} \Gamma^{\delta a}_{d]b} + 2\Gamma^{\delta a}_{q[c} \Gamma^{\delta q}_{d]b}$$


Metric spacetime geometry:

- fundamental tuple (M , $g(x)$)
spacetime with metric
- the lifted Levi-Civita connection $p\Gamma$
determines the geometry of phase space
- the Levi-Civita connection and
the Riemann curvature determine
the geometry of spacetime

$$\Gamma^a_{bc} = \frac{1}{2} g^{aq} (\partial_b g_{cq} + \partial_c g_{bq} - \partial_q g_{bc})$$

$$\delta_a = \partial_a - p_c \Gamma^c_{ab}(x) \bar{\partial}^b$$

$$R^a_{bcd}(x) = 2\partial_{[c} \Gamma^a_{d]b} + 2\Gamma^a_{q[c} \Gamma^q_{d]b}$$



Hamilton Geometry

Hamilton phase space geometry

- fundamental tuple (T^*M , $H(x,p)$)
cotangent bundle with Hamiltonian
- the Cartan non-linear connection N
determines the geometry of phase space
- the spacetime connection and its curvature
determine the geometry of spacetime
- the momentum space connection
and its curvature determine the
geometry of momentum space

$$C^{bc}{}_a = \frac{1}{2} g_{aq}^H \bar{\partial}^b g^{Hbq}$$


$$Q_a{}^{bcd}(x, p) = 2\bar{\partial}^{[c} C^{d]b}{}_a + 2C^{q[c}{}_a C^{d]b}{}_q$$

Metric spacetime geometry:

- fundamental tuple (M , $g(x)$)
spacetime with metric
- the lifted Levi-Civita connection $p\Gamma$
determines the geometry of phase space
- the Levi-Civita connection and the Riemann
curvature determine the geometry of spacetime
- the momentum space connection
and its curvature vanish,
momentum space is flat

$$C^{bc}{}_a = 0$$

$$Q_a{}^{bcd}(x, p) = 0$$



Hamilton Geometry

Hamilton phase space geometry

- fundamental tuple ($T^*M, H(x,p)$)
cotangent bundle with Hamiltonian
- the Cartan non-linear connection N
determines the geometry of phase space
- the spacetime connection and its curvature
determine the geometry of spacetime
- the momentum space connection and its
curvature determine the geometry of momentum
space

Metric spacetime geometry:

- fundamental tuple ($M, g(x)$)
spacetime with metric
- the lifted Levi-Civita connection $p\Gamma$
determines the geometry of phase space
- the Levi-Civita connection and the Riemann
curvature determine the geometry of spacetime
- the momentum space connection and its
curvature vanish, momentum space is flat

The geometry of a first order modified dispersion relation

[L. Barcaroli, L. Brunkhorst, G. Gubitosi, N. Loret, CP 2015]

$$H = g^{ab}(x)p_ap_b + \ell G^{abc}(x)p_ap_bp_c$$

$$N_{ab}(x, p) = -\Gamma^q_{ab}p_q + \ell \frac{3}{4}p_qp_r(\nabla_a G^{qr}_b + \nabla_b G^{qr}_a - g^{sq}\nabla_s G^r_{ab}) + \mathcal{O}(\ell^2)$$

$$\Gamma^{\delta a}_{bc}(x, p) = \Gamma^a_{bc} + \ell \frac{2}{3}p_q g^{ad}(\nabla_d G_{bc}^q - \nabla_b G_{cd}^q - \nabla_c G_{bd}^q) + \mathcal{O}(\ell^2)$$

$$C^{bc}_a(x, p) = \ell \frac{3}{2}G^{ab}_c + \mathcal{O}(\ell^2).$$

Hamilton Geometry

Hamilton phase space geometry

- fundamental tuple ($T^*M, H(x,p)$)
cotangent bundle with Hamiltonian
- the Cartan non-linear connection N
determines the geometry of phase space
- the spacetime connection and its curvature
determine the geometry of spacetime
- the momentum space connection and its
curvature determine the geometry of momentum
space

Metric spacetime geometry:

- fundamental tuple ($M, g(x)$)
spacetime with metric
- the lifted Levi-Civita connection $p\Gamma$
determines the geometry of phase space
- the Levi-Civita connection and the Riemann
curvature determine the geometry of spacetime
- the momentum space connection and its
curvature vanish, momentum space is flat

The geometry of a first order modified dispersion relation

$$H = g^{ab}(x)p_ap_b + \ell G^{abc}(x)p_ap_bp_c$$

$$N_{ab}(x, p) = -\Gamma^q_{ab}p_q + \ell \frac{3}{4}p_qp_r(\nabla_a G^{qr}_b + \nabla_b G^{qr}_a - g^{sq}\nabla_s G^r_{ab}) + \mathcal{O}(\ell^2)$$

$$\Gamma^{\delta a}_{bc}(x, p) = \Gamma^a_{bc} + \ell \frac{2}{3}p_q g^{ad}(\nabla_d G_{bc}^q - \nabla_b G_{cd}^q - \nabla_c G_{bd}^q) + \mathcal{O}(\ell^2)$$

$$C^{bc}_a(x, p) = \ell \frac{3}{2}G^{ab}_c + \mathcal{O}(\ell^2).$$

The geometry of phase space is determined by three distinguished objects

- the Cartan non-linear connection
 - the spacetime connection and its curvature
 - the momentum space connection and its curvature
- all derived canonically from a Hamiltonian/disp. relation on phase space.

Geometric interpretation of Hamilton equations of motion

Point particle follow solutions of Hamilton equations of motion

$$\dot{p}_a + \partial_a H = 0, \quad \dot{x}^a = \bar{\partial}^a H$$

In terms of the non-linear connection N_{ab}

$$\dot{p}_a + N_{ab} \bar{\partial}^b H = -[\partial_a - N_{ab} \bar{\partial}^b] H = -\delta_a H, \quad \dot{x}^a = \bar{\partial}^a H.$$

They can be compared to the autoparallels of the geometry

$$\dot{p}_a + N_{ab} \bar{\partial}^b H = 0.$$

In general point particles are not freely-falling but subject to a force term

$$\delta_a H$$

Theorem: [L. Barcaroli, L. Brunkhorst, G. Gubitosi, N. Loret, CP 2015]

For homogeneous Hamiltonians $H(x, s, p) = s^r H(x, p)$
solutions to the Hamilton equations of motion are autoparallels:
the source term vanishes

$$\delta_a H = 0$$

1. Dispersion Relations and the Geometry of Spacetime
2. Hamiltonian Phase Space Geometry
3. Conclusion and Outlook

Summary

Dispersion relations are level sets of Hamilton functions on phase space.

The geometry of phase space is determined by three distinguished objects

- the Cartan non-linear connection
- the spacetime connection and its curvature
- the momentum space connection and its curvature

The spacetime and momentum space geometry is phase space dependent, i.e. depend on positions x and momenta p .

An only x dependent geometry of spacetime and only p dependent geometry of momentum space is very special.

Summary

Dispersion relations are level sets of Hamilton functions on phase space.

The geometry of phase space is determined by three distinguished objects

- the Cartan non-linear connection
- the spacetime connection and its curvature
- the momentum space connection and its curvature

The spacetime and momentum space geometry is phase space dependent, i.e. depend on positions x and momenta p .

An only x dependent geometry of spacetime and only p dependent geometry of momentum space is very special.

The geometric framework of Hamilton geometry allows a precise geometric comparison between general relativity and the geometry of spacetime induced by general (modified) dispersion relations.

Outlook - Hamilton Geometry

Quantum gravity phenomenology:

- Geometric understanding of (modified) addition of momenta
- Geometric definition of observers, and transformations between them
- Study modified dispersion relation in homogeneous and isotropic, spherically symmetric or axially symmetric spacetime geometries

Applications to the analysis of PDEs:

- Dispersion relations are local representations of PDEs
- Geometric description of propagation of field modes.

Example: Propagation of light in general linear electrodynamics, for example in media

$$H(x, p) = \mathcal{G}^{abcd}(x) p_a p_b p_c p_d$$

Dispersion relations are level sets of Hamilton functions:
They determine the geometry of spacetime and momentum space.

Phys. Rev. D 92 (2015) 8, 084053; arXiv:1507.00922:

Hamilton geometry: Phase space geometry from modified dispersion relations

Thank you for your attention