

Black hole non-modal linear stability: the $\Lambda \geq 0$ Schwarzschild case

talk based on PRL **112** (2014) 191101 and arXiv 1603.03749

Gustavo Dotti

FaMAF, Universidad Nacional de Córdoba
ARGENTINA

GR21, Columbia University
July 13th, 2016

Dedicated to Sergio Dain

Defining linear stability

Linear vacuum Einstein equation with cosmological constant (LEE)

$$\mathcal{E}[\delta g_{\alpha\beta}] := -\frac{1}{2}\nabla^\gamma\nabla_\gamma\delta g_{\alpha\beta} - \frac{1}{2}\nabla_\alpha\nabla_\beta(g^{\gamma\delta}\delta g_{\gamma\delta}) + \nabla^\gamma\nabla_{(\alpha}\delta g_{\beta)\gamma} - \Lambda\delta g_{\alpha\beta} = 0$$

We are interested in equivalence classes $[\delta g_{\alpha\beta}]$ of slns mod gauge transformations:

$$\delta g_{\alpha\beta} \sim \delta g_{\alpha\beta} + \nabla_\alpha\xi_\beta + \nabla_\beta\xi_\alpha$$

Rough stability idea: The outer region of a stationary black hole is linearly stable if linear metric perturbations *do not grow unbounded*.

Expectation: *perturbed Kerr BH metrics decay to a nearby Kerr BHs.*

Defining linear stability

Linear vacuum Einstein equation with cosmological constant (LEE)

$$\mathcal{E}[\delta g_{\alpha\beta}] := -\frac{1}{2}\nabla^\gamma\nabla_\gamma\delta g_{\alpha\beta} - \frac{1}{2}\nabla_\alpha\nabla_\beta(g^{\gamma\delta}\delta g_{\gamma\delta}) + \nabla^\gamma\nabla_{(\alpha}\delta g_{\beta)\gamma} - \Lambda\delta g_{\alpha\beta} = 0$$

We are interested in equivalence classes $[\delta g_{\alpha\beta}]$ of slns mod gauge transformations:


$$\delta g_{\alpha\beta} \sim \delta g_{\alpha\beta} + \nabla_\alpha\xi_\beta + \nabla_\beta\xi_\alpha$$

Rough stability idea: The outer region of a stationary black hole is linearly stable if linear metric perturbations *do not grow unboundedly*.


Expectation: *perturbed Kerr BH metrics decay to a nearby Kerr BHs.*

- If $\Lambda \geq 0$ both II and $II' \cup I' \cup I'$ are globally hyperbolic.
- If $\Lambda < 0$ boundary conditions have to be specified at the conformal timelike boundary and there are instabilities if some Robin boundary conditions are chosen.

Carter-Penrose diagram: II outer static region; I : inner non static region



Zero or positive cosmological constant



Negative cosmological constant

Modal approach: i) solving the LEE

$$\underbrace{g_{\alpha\beta} dz^\alpha dz^\beta}_{\nabla_\alpha, \epsilon_{\alpha\beta\gamma\delta}} = \underbrace{g_{ab}(x) dx^a dx^b}_{\text{orbit space: } D_C, \epsilon_{ab}} + r^2(x) \underbrace{\hat{g}_{ij}(y) dy^i dy^j}_{S^2: \hat{D}_K, \epsilon_{ij}}$$

$$= -f(r) dt^2 + \frac{dr^2}{f(r)} + r^2 (d\theta^2 + \sin^2(\theta) d\phi^2)$$

$$\mathbf{J}^2 = \sum_{k=1}^3 (\mathbf{J}_k)^2, \quad (\mathbf{J}_3 = \partial_\phi, \text{ etc}) \quad \text{and} \quad P : (t, r, \theta, \phi) \rightarrow (t, r, \pi - \theta, \phi + \pi)$$

$$\delta g_{\alpha\beta} = \sum_{p=\pm, \ell, m} \delta g_{\alpha\beta}^{(\ell, m, p)} \quad \text{where} \quad \mathbf{J}^2 \delta g_{\alpha\beta}^{(\ell, m, \pm)} = -\ell(\ell+1) \delta g_{\alpha\beta}^{(\ell, m, \pm)}$$

$$\text{even (+) and odd (-) modes} \quad P_* \delta g_{\alpha\beta}^{(\ell, m, \pm)} = \pm (-1)^\ell \delta g_{\alpha\beta}^{(\ell, m, \pm)}$$

$$\ell \geq 2 \text{ LEE equivalent to infinite set of 1+1 wave eqns: } \underbrace{g^{ab} D_a D_b \phi_{(\ell, m)}^\pm - U_\ell^\pm \phi_{(\ell, m)}^\pm}_{\text{LEE for } \ell \geq 2 \text{ modes}} = 0$$

Modal approach: i) solving the LEE

	odd modes ($p = -$)	even modes ($p = +$)
$\ell = 0$		δM shift within Kerr family
$\ell = 1$	$\delta j^{(k)}$ shift within Kerr family	
$\ell \geq 2$	$\delta g_{\alpha\beta}^{(\ell,m,-)} = \mathcal{D}_{\alpha\beta}^{(-)}[\phi_{(\ell,m)}^-, S_{(\ell,m)}]$	$\delta g_{\alpha\beta}^{(\ell,m,+)} = \mathcal{D}_{\alpha\beta}^{(+)}[\phi_{(\ell,m)}^+, S_{(\ell,m)}]$

$S_{(\ell,m)}$ are spherical harmonics: $\hat{g}^{ij}\hat{D}_i\hat{D}_j S_{(\ell,m)} = -\ell(\ell+1)S_{(\ell,m)}$,

$\phi_{(\ell,m)}^\pm$ are solutions of 2D wave eqns: $\underbrace{g^{ab}D_a D_b \phi_{(\ell,m)}^\pm - U_\ell^\pm \phi_{(\ell,m)}^\pm}_{\text{LEE for } \ell \geq 2 \text{ modes}} = 0$

Generic perturbations parametrized by gauge invariant fields and constants: $\delta M, \delta j^{(k)}$ (constants read from initial datum) and $\phi_{(\ell,m)}^\pm$ (dynamical fields)

Odd sector non modal approach: linear stability

- Standard approach to stability problem consists on setting bounds on the infinite set of fields $\phi_{(\ell,m)}(t, r)$
- Up to 2 derivatives of these fields enter $\delta g_{\alpha\beta}$, thus 4 derivatives in $\delta R^\alpha{}_{\beta\gamma\delta}$.
- Any geometric implies $\sum_{(\ell,m)}$
- Implications of the boundedness (either integral or pointwise) of the $\phi_{(\ell,m)}^\pm$ on the perturbed geometry not obvious a priori
- Look to parametrize the space \mathcal{L} of linearized solutions of the Einstein's equations around Schwarzschild de Sitter BH with geometrical fields as an alternative to

$$\mathcal{L} = \{\delta M, \delta j^{(k)}, \phi_{(\ell,m)}^\pm\}$$

- Need to estimate the growth of these fields in order to analyze stability

RW equation and 4D-RW equation:

$$\begin{aligned} \ell \geq 2 \text{ LEE} &\Leftrightarrow \overbrace{\left[\partial_t^2 - \partial_{r^*}^2 + f \left(\frac{\ell(\ell+1)}{r^2} - \frac{6M}{r^3} \right) \right] \phi_{(\ell,m)}^- = 0, \quad \hat{D}^k \hat{D}_k S_{(\ell,m)} = -\ell(\ell+1) S_{(\ell,m)}}^{\text{RW equation}} \\ &\Leftrightarrow \underbrace{\left[\nabla_\alpha \nabla^\alpha + \frac{8M}{r^3} - \frac{2\Lambda}{3} \right] \Phi = 0}_{\text{4D RW equation}} \quad \Phi = \sum_{(\ell \geq 2, m)} \frac{\phi_{(\ell,m)}^-}{r} S_{(\ell,m)} \end{aligned}$$

Odd sector non modal approach: perturbed curvature scalars

We consider the effect of a perturbation on the curvature scalars:

$$Q_- = \frac{1}{48} {}^* C^{\alpha\beta\gamma\delta} C_{\alpha\beta\gamma\delta}, \quad Q_+ = \frac{1}{48} C^{\alpha\beta\gamma\delta} C_{\alpha\beta\gamma\delta}, \quad X = \frac{1}{720} (\nabla_\epsilon C_{\alpha\beta\gamma\delta}) (\nabla^\epsilon C^{\alpha\beta\gamma\delta}).$$

The background values are:

$$Q_{-S(A)dS} = 0, \quad Q_{+S(A)dS} = \frac{M^2}{r^6}, \quad X_{S(A)dS} = \frac{M^2}{r^9} (r - 2M) - \frac{\Lambda M^2}{3r^6}$$

This implies that the following fields are gauge invariant:

$$G_- = \delta Q_- \quad \text{and} \quad G_+ = (9M - 4r + \Lambda r^3) \delta Q_+ + 3r^3 \delta X$$

$G_- = G_-[\delta g^{(-)}]$, this functional depends on up to four derivatives of the $\phi_{(\ell,m)}^-$.

Using repeatedly the LEE we arrive at: (recall that $\mathcal{L} = \{\overbrace{\delta j^{(k)}}^{\text{odd}}, \overbrace{\phi_{(\ell,m)}^-, \delta M, \phi_{(\ell,m)}^+}^{\text{even}}\}$)

$$G_- = -\frac{6M}{r^7} \sqrt{\frac{4\pi}{3}} \sum_{m=1}^3 \delta j^{(m)} S_{(1,m)} - \underbrace{\frac{3M}{r^5} \sum_{(\ell \geq 2, m)} \frac{(\ell+2)!}{(\ell-2)!} \frac{\phi_{(\ell,m)}^-}{r} S_{(\ell,m)}}_{J^2(J^2+2)\Phi}$$

Odd sector non modal approach: linear stability

$$G_- = -\frac{6M}{r^7} \sqrt{\frac{4\pi}{3}} \sum_{m=1}^3 \delta j^{(m)} S_{(1,m)} - \frac{3M}{r^5} \sum_{(\ell \geq 2, m)} \frac{(\ell+2)!}{(\ell-2)!} \frac{\phi_{(\ell,m)}^-}{r} S_{(\ell,m)}$$

From G_- we can recover $(\delta j^{(m)}, \phi_{(\ell,m)}^-)$ and therefore $\delta g_{\alpha\beta}^{(-)}$ in a given gauge

All the gauge invariant information in $\delta g_{\alpha\beta}^{(-)}$ is encoded in G_-

$[\delta g_{\alpha\beta}^{(-)}] \rightarrow G_- \left([\delta g_{\alpha\beta}^{(-)}] \right)$ is a bijection

Odd LEE are entirely equivalent to $\left[\nabla_\alpha \nabla^\alpha + \frac{8M}{r^3} - \frac{2\Lambda}{3} \right] (r^5 G_-) = 0!!!$ (4DRWE)

Boundedness: For any smooth solution of the odd LEE which has compact support on Cauchy surfaces of the extended $I \cup II \cup I' \cup II'$ Schwarzschild (Schwarzschild de Sitter) BH, there exists a constant K_- such that $|G_-| < K_- r^{-6}$ for $r > r_h$ ($r_h < r < r_c$)

Decay: For large t Price/Brady tails give a slowly rotating Kerr/dS BH:

$$G_- \simeq -\frac{6M}{r^7} \sqrt{\frac{4\pi}{3}} \sum_{m=1}^3 \delta j^{(m)} S_{(1,m)}$$

Even perturbations: difficulties for a 4D approach

1) For odd perturbations

$$\begin{aligned} & \mathcal{H}_\ell^- \text{ (RW Hamiltonian)} \\ \ell \geq 2 \text{ odd LEE} & \Leftrightarrow \overbrace{[\partial_t^2 - \partial_{r^*}^2 + f \left(\frac{\ell(\ell+1)}{r^2} - \frac{6M}{r^3} \right)]} \phi_{(\ell,m)}^- = 0, \quad \hat{D}^k \hat{D}_k S_{(\ell,m)} = -\ell(\ell+1) S_{(\ell,m)} \\ & \Leftrightarrow \underbrace{\left[\nabla_\alpha \nabla^\alpha + \frac{8M}{r^3} - \frac{2\Lambda}{3} \right]}_{\text{4D RW equation}} \Phi = 0 \quad \Phi = \sum_{(\ell \geq 2, m)} \frac{\phi_{(\ell,m)}^-}{r} S_{(\ell,m)} \end{aligned}$$

2) For even perturbations the 2D Zerilli wave equation is not related to a 4D equation:

$$\mathcal{H}_\ell^+ = -\partial_{r^*}^2 + f V_{(\ell,m)}^Z \quad (\text{Zerilli Hamiltonian})$$

with potential ($\mu = (\ell - 1)(\ell + 2)$)

$$V_{(\ell,m)}^Z = \frac{[\mu^2 \ell(\ell + 1) - 24M^2 \Lambda] r^3 + 6\mu^2 M r^2 + 36\mu M^2 r + 72M^3}{r^3 (6M + ((\ell - 1)(\ell + 2))^2 r^2)}.$$

Even perturbations: difficulties for a 4D approach

3) For odd perturbations there is a gauge invariant scalar related to the perturbed Weyl tensor (thus expected to satisfy some kind of wave equation)

$$G_- = \delta Q_- = \delta \left(\frac{1}{48} {}^* C^{\alpha\beta\gamma\delta} C_{\alpha\beta\gamma\delta} \right)$$

4) For even perturbations we need to use differential invariants to construct curvature related gauge invariant perturbation fields. The simplest such field is

$$G_+ = G_+ = (9M - 4r + \Lambda r^3) \delta Q_+ + 3r^3 \delta X$$

$$Q_+ = \frac{1}{48} C^{\alpha\beta\gamma\delta} C_{\alpha\beta\gamma\delta}, \quad X = \frac{1}{720} (\nabla_\epsilon C_{\alpha\beta\gamma\delta}) (\nabla^\epsilon C^{\alpha\beta\gamma\delta}).$$

These are not expected to satisfy wave equations as a consequence of the LEE !

Even perturbations: Chandrasekhar's even \leftrightarrow odd duality

Chandrasekhar (80's) found that

$$\mathcal{H}_\ell^\pm = \mathcal{D}_\ell^\pm \mathcal{D}_\ell^\mp - E_\ell^2, \quad \mathcal{D}_\ell^\pm = \pm \frac{\partial}{\partial r^*} + W_\ell(r)$$

\mathcal{D}_ℓ^+ maps solutions of the odd (RW) equations to solutions of the even (Z) equation:

$$(\partial_t^2 + \mathcal{H}_\ell^-) \phi_{(\ell,m)}^- = 0 \Rightarrow (\partial_t^2 + \mathcal{H}_\ell^+) (\mathcal{D}_\ell^+ \phi_{(\ell,m)}^-) = 0$$

- The general solution of the differential equation $\mathcal{D}_\ell^+ \chi = 0$ is a constant times

$$\chi_\ell^- = \frac{(\ell+2)(\ell-1)r + 6M}{r} \exp(-w_\ell r^*) \notin L^2(\mathbb{R}, dr^*)$$

- This implies that $\mathcal{D}_\ell^+ : L^2(\mathbb{R}, dr^*) \rightarrow L^2(\mathbb{R}, dr^*)$ is injective (also in more general spaces, since the above sln blows up at the BS).

Assume $\Lambda \geq 0$. For any solution $\phi_{(\ell,m)}^+$ of the ZE in $L^2(\mathbb{R}_{r^*}, dr^*)$ there is a unique solution $\phi_{(\ell,m)}^-$ of RWE in $L^2(\mathbb{R}_{r^*}, dr^*)$ such that $\phi_{(\ell,m)}^+ = \mathcal{D}_\ell^+ \phi_{(\ell,m)}^-$.

Even perturbations: Non modal stability

$$G_+ = -\frac{2M\delta M}{r^5} + \frac{M}{2r^4} \sum_{\ell \geq 2} \frac{(\ell+2)!}{(\ell-2)!} [f\partial_r + Z_\ell] \phi_{(\ell,m)}^+ S_{(\ell,m)}, \quad Z_\ell = Z_\ell(r)$$

From G_+ we can recover $(\delta M, \phi_{(\ell,m)}^+)$ and therefore $\delta g_{\alpha\beta}^{(+)}$ in a given gauge

All the gauge invariant information in $\delta g_{\alpha\beta}^{(+)}$ is encoded in G_+

$[\delta g_{\alpha\beta}^{(+)}] \rightarrow G_+ \left([\delta g_{\alpha\beta}^{(+)}] \right)$ is a bijection

$$G_+ = -\frac{2M\dot{M}}{r^5} + \frac{M}{2r^3} \left[\partial_t^2 \Phi_5 + \sum_{j=0}^4 r^{-j} \Phi_j + \frac{f(r-3M)}{r^3} \Phi_5 + \frac{f}{r} \Phi_6 \right] + \frac{M}{2r^3} \partial_{r^*} \Phi_6 + \frac{M(r-3M)}{2r^5} \partial_{r^*} \Phi_5,$$

(using Chandra's duality) $\Phi_j = \sum_{(\ell \geq 2, m)} P_j(\ell) \frac{\phi_{(\ell,m)}^-}{r} S_{(\ell,m)}, \quad j = 0, \dots, 6$ satisfy the 4DRWE !!!

Boundedness: For any smooth solution of the odd LEE which has compact support on Cauchy surfaces of the extended $I \cup II \cup I' \cup II'$ Schwarzschild (or Schwarzschild de Sitter) BH, there exists a constant K_+ such that $|G_+| < K_+ r^{-3}$ for $r > r_h$ ($r_h < r < r_c$)

- All gauge invariant perturbation info contained in scalar curvature perturbation fields

$$G_- = -\frac{6M}{r^7} \sqrt{\frac{4\pi}{3}} \sum_{m=1}^3 \delta j^{(m)} S_{(1,m)} - \frac{3M}{r^5} \sum_{(\ell \geq 2, m)} \frac{(\ell+2)!}{(\ell-2)!} \frac{\phi_{(\ell,m)}^-}{r} S_{(\ell,m)}$$

$$G_+ = -\frac{2M\delta M}{r^5} + \frac{M}{2r^4} \sum_{\ell \geq 2} \frac{(\ell+2)!}{(\ell-2)!} [f\partial_r + Z_\ell] \phi_{(\ell,m)}^+ S_{(\ell,m)}$$

- $[\delta g_{\alpha\beta}] \rightarrow (G_-([\delta g_{\alpha\beta}]), G_+([\delta g_{\alpha\beta}]))$ is a bijection
- LEE entirely equivalent to 4D RW equation !

Summary

- All gauge invariant perturbation info contained in scalar curvature perturbation fields

$$G_- = -\frac{6M}{r^7} \sqrt{\frac{4\pi}{3}} \sum_{m=1}^3 \delta j^{(m)} S_{(1,m)} - \frac{3M}{r^5} \sum_{(\ell \geq 2, m)} \frac{(\ell+2)!}{(\ell-2)!} \frac{\phi_{(\ell,m)}^-}{r} S_{(\ell,m)}$$
$$G_+ = -\frac{2M\delta M}{r^5} + \frac{M}{2r^4} \sum_{\ell \geq 2} \frac{(\ell+2)!}{(\ell-2)!} [f\partial_r + Z_\ell] \phi_{(\ell,m)}^+ S_{(\ell,m)}$$

- $[\delta g_{\alpha\beta}] \rightarrow (G_-([\delta g_{\alpha\beta}]), G_+([\delta g_{\alpha\beta}]))$ is a bijection
- LEE entirely equivalent to 4D RW equation !
- Perturbations are bounded in the outer region:

$$|G_-| \leq K_-/r^6, \quad |G_+| \leq K_+/r^3$$

Summary

- All gauge invariant perturbation info contained in scalar curvature perturbation fields

$$G_- = -\frac{6M}{r^7} \sqrt{\frac{4\pi}{3}} \sum_{m=1}^3 \delta j^{(m)} S_{(1,m)} - \frac{3M}{r^5} \sum_{(\ell \geq 2, m)} \frac{(\ell+2)!}{(\ell-2)!} \frac{\phi_{(\ell,m)}^-}{r} S_{(\ell,m)}$$

$$G_+ = -\frac{2M\delta M}{r^5} + \frac{M}{2r^4} \sum_{\ell \geq 2} \frac{(\ell+2)!}{(\ell-2)!} [f\partial_r + Z_\ell] \phi_{(\ell,m)}^+ S_{(\ell,m)}$$

- $[\delta g_{\alpha\beta}] \rightarrow (G_-([\delta g_{\alpha\beta}]), G_+([\delta g_{\alpha\beta}]))$ is a bijection
- LEE entirely equivalent to 4D RW equation !
- Perturbations are bounded in the outer region:

$$|G_-| \leq K_-/r^6, \quad |G_+| \leq K_+/r^3$$

- For large t we get decay within Kerr/ds family:

$$G_- \simeq -\frac{6M}{r^7} \sqrt{\frac{4\pi}{3}} \sum_{m=1}^3 \delta j^{(m)} S_{(1,m)}, \quad G_+ \simeq -\frac{2M\delta M}{r^5}$$