


# Localised $\text{AdS}_5 \times S^5$ Black Holes

Jorge E. Santos

GR 21 - New York



In collaboration with  
Óscar J. C. Dias and Benson Way

# The original AdS/CFT - '97 Maldacena

Where it all began:

**Type IIB supergravity theory** on  $\text{AdS}_5 \times S^5$  with radius  $L$  and  $N$  units of  $F_{(5)}$  flux on  $S^5$



The large  $N$  limit and strong coupling limit of  $\mathcal{N} = 4$  **Super Yang-Mills (SYM) theory** with t'Hooft coupling  $\lambda$ .


# The original AdS/CFT - '97 Maldacena

Where it all began:

**Type IIB supergravity theory** on  $\text{AdS}_5 \times S^5$  with radius  $L$  and  $N$  units of  $F_{(5)}$  flux on  $S^5$



The large  $N$  limit and strong coupling limit of  $\mathcal{N} = 4$  **Super Yang-Mills (SYM) theory** with t'Hooft coupling  $\lambda$ .



$\mathcal{N} = 4$  SYM on Einstein static universe - '98 Witten

## The role of Black Holes:

- Black holes radiate at **temperature**  $T$  - '74 Hawking.

# $\mathcal{N} = 4$ SYM on Einstein static universe - '98 Witten

## The role of Black Holes:

- Black holes radiate at **temperature**  $T$  - '74 Hawking.
- Witten **conjectured** that black holes in global  $\text{AdS}_5$  should be **dual to the hot** phase of  $\mathcal{N} = 4$  SYM on  $\mathbb{R}_t \times S^3_{\mathbf{L}}$ .

# $\mathcal{N} = 4$ SYM on Einstein static universe - '98 Witten

## The role of Black Holes:

- Black holes radiate at **temperature**  $T$  - '74 Hawking.
- Witten **conjectured** that black holes in global  $\text{AdS}_5$  should be **dual to the hot** phase of  $\mathcal{N} = 4$  SYM on  $\mathbb{R}_t \times S^3_{\mathbf{L}}$ .
- Black holes should be the **the holographic dual to the end state of a real time** thermalisation process - **maximum entropy at fixed energy**.

# $\mathcal{N} = 4$ SYM on Einstein static universe - '98 Witten

## The role of Black Holes:

- Black holes radiate at **temperature**  $T$  - '74 Hawking.
- Witten **conjectured** that black holes in global  $\text{AdS}_5$  should be **dual to the hot** phase of  $\mathcal{N} = 4$  SYM on  $\mathbb{R}_t \times S^3_{\text{L}}$ .
- Black holes should be the **the holographic dual to the end state of a real time** thermalisation process - **maximum entropy at fixed energy**.

## Question:

What is the phase diagram of static black holes in global  $\text{AdS}_5 \times S^5$ ?

# The equations of motion

Keeping it *simple*:

- Type IIB supergravity has many fields:



# The equations of motion

Keeping it *simple*:

- Type IIB supergravity has many fields:
  - one symmetric two tensor -  $g$
  - one real self dual five-form -  $F$
  - one complex three-form -  $G_3$
  - one complex scalar -  $B$

# The equations of motion

Keeping it *simple*:

- Type IIB supergravity has many fields:
  - **one symmetric two tensor** -  $g$
  - **one real self dual five-form** -  $F$
  - **one complex three-form** -  $G_3$
  - **one complex scalar** -  $B$
- Can consistently set  $G_3 = 0$  and  $B = 1$ .

# The equations of motion

Keeping it *simple*:

- Type IIB supergravity has many fields:
  - **one symmetric two tensor** -  $g$
  - **one real self dual five-form** -  $F$
  - **one complex three-form** -  $G_3$
  - **one complex scalar** -  $B$
- Can consistently set  $G_3 = 0$  and  $B = 1$ .

Reduced equations of motion:

$$R_{ab} = \frac{1}{48} F_a{}^{cdef} F_{bcdef}, \quad \nabla_a F^{abcde} = 0 \quad \text{and} \quad \star F = F.$$

# The smeared black hole

A smeared family of solutions - '80 Freund & Rubin:

**Any** solution to:

$$R_{\mu\nu}^{(5)} = \frac{4}{\mathbf{L}^2} g_{\mu\nu}^{(5)}$$

can be **oxidised** to **ten-dimensions** via

$$ds^2 = g_{\mu\nu}^{(5)} dx^\mu dx^\nu + \mathbf{L}^2 d\Omega_5^2 \quad \text{and} \quad F = \text{Vol}_{\text{AdS}_5} + \text{Vol}_{S^5} .$$

# The smeared black hole

## A smeared Black Hole:

The **Schwarzschild-AdS<sub>5</sub>** black hole solution lifts to:

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega_3^2 + \mathbf{L}^2 d\Omega_5^2,$$

where

$$f(r) = \frac{r^2}{\mathbf{L}^2} + 1 - \frac{r_+^2}{r^2} \left( \frac{r_+^2}{\mathbf{L}^2} + 1 \right),$$

with  $y_+ \equiv r_+/\mathbf{L}$  measuring the **black hole radius** with respect to the **radius of the S<sup>5</sup>** (also the radius of the S<sup>3</sup> at  $\partial$ ).

# The smeared black hole

## A smeared Black Hole:

The **Schwarzschild- $\text{AdS}_5$**  black hole solution lifts to:

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega_3^2 + \mathbf{L}^2 d\Omega_5^2,$$

where


$$f(r) = \frac{r^2}{\mathbf{L}^2} + 1 - \frac{r_+^2}{r^2} \left( \frac{r_+^2}{\mathbf{L}^2} + 1 \right),$$

with  $y_+ \equiv r_+/\mathbf{L}$  measuring the **black hole radius** with respect to the **radius of the  $S^5$**  (also the radius of the  $S^3$  at  $\partial$ ).

## Thermodynamic Properties:

$$E = \frac{3\mathbf{N}^2}{4\mathbf{L}} y_+^2 (1 + y_+^2), \quad S = \pi y_+^2 \mathbf{N}^2 \quad \text{and} \quad T = \frac{1 + 2y_+^2}{2\pi y_+ \mathbf{L}}.$$

# Microcanonical Ensemble



Key question:

Is this the unique static solution with the desired asymptotics?


## Size Matters - '98 Banks *et al.*

When black hole radius **is small**, i.e.  $y_+ \ll 1$ :




## Size Matters - '98 Banks *et al.*

When black hole radius **is small**, i.e.  $y_+ \ll 1$ :



The separation of **horizon length scales** suggests a  
**Gregory-Laflamme instability**:  
heuristics confirmed '02 Hubeny & Rangamani.

# Instability Growth Rate - '15 Buchel, Lehner




Alike in **Gregory-Laflamme instability**, **zero-mode** -  $y_+ \sim 0.44$   
 - connects to novel **non-uniform solutions** along the  $S^5$  - with  
 horizon topology  $S^3 \times S^5$ .

Lumpy holes in  $\text{AdS}_5 \times S^5$  - '15 Dias, Santos and Way

Lumpy black holes:


- **Lumpy holes** have a **lower entropy** than smeared Schwarzschild- $\text{AdS}_5$  at **fixed energy**.



# Lumpy holes in $\text{AdS}_5 \times S^5$ - '15 Dias, Santos and Way

## Lumpy black holes:

- **Lumpy holes** have a **lower entropy** than smeared Schwarzschild- $\text{AdS}_5$  at **fixed energy**.



- They cannot be the **endpoint of the instability**!

## Other candidate endpoints - '98 Banks *et al.*

### Localised black holes:

- At least one other candidate endpoint exists: a **localised black hole in  $\text{AdS}_5 \times S^5$** .

## Other candidate endpoints - '98 Banks *et al.*

### Localised black holes:

- At least one other candidate endpoint exists: a **localised black hole in  $\text{AdS}_5 \times S^5$** .
- Must have the following property: **higher entropy** than smeared Schwarzschild- $\text{AdS}_5$  hole, at fixed energy.

## Other candidate endpoints - '98 Banks *et al.*

### Localised black holes:

- At least one other candidate endpoint exists: a **localised black hole in  $\text{AdS}_5 \times S^5$** .
- Must have the following property: **higher entropy** than smeared Schwarzschild- $\text{AdS}_5$  hole, at fixed energy.
- If true, this would imply a **violation of weak cosmic censorship** in  $\text{AdS}_5 \times S^5$ .

## Other candidate endpoints - '98 Banks *et al.*

### Localised black holes:

- At least one other candidate endpoint exists: a **localised black hole in  $\text{AdS}_5 \times S^5$** .
- Must have the following property: **higher entropy** than smeared Schwarzschild- $\text{AdS}_5$  hole, at fixed energy.
- If true, this would imply a **violation of weak cosmic censorship** in  $\text{AdS}_5 \times S^5$ .

### Vanilla flavour Gregory-Laflamme instability:



## Other candidate endpoints - '98 Banks *et al.*

### Localised black holes:

- At least one other candidate endpoint exists: a **localised black hole in  $\text{AdS}_5 \times S^5$** .
- Must have the following property: **higher entropy** than smeared Schwarzschild- $\text{AdS}_5$  hole, at fixed energy.
- If true, this would imply a **violation of weak cosmic censorship** in  $\text{AdS}_5 \times S^5$ .

### Vanilla flavour Gregory-Laflamme instability:

- For the **original Gregory-Laflamme instability**, this is indeed what happens: '10 Pretorius and Lehner.

## Other candidate endpoints - '98 Banks *et al.*

### Localised black holes:

- At least one other candidate endpoint exists: a **localised black hole in  $\text{AdS}_5 \times S^5$** .
- Must have the following property: **higher entropy** than smeared Schwarzschild- $\text{AdS}_5$  hole, at fixed energy.
- If true, this would imply a **violation of weak cosmic censorship** in  $\text{AdS}_5 \times S^5$ .

### Vanilla flavour Gregory-Laflamme instability:

- For the **original Gregory-Laflamme instability**, this is indeed what happens: '10 Pretorius and Lehner.
- **Localised solutions** were first constructed in '05 by Kudoh and Wiseman.

# Localised Black Holes in $\text{AdS}_5 \times S^5$

- These **localised black holes** have **horizon topology**  $S^8$  and asymptote to  $\text{AdS}_5 \times S^5$ , whose **constant time and 'holographic' slices** have topology  $S^3 \times S^5$ .

# Localised Black Holes in $\text{AdS}_5 \times S^5$

- These **localised black holes** have **horizon topology**  $S^8$  and asymptote to  $\text{AdS}_5 \times S^5$ , whose **constant time and 'holographic' slices** have topology  $S^3 \times S^5$ .
- Can think of it as **'inverted' black rings**.

# Localised Black Holes in $\text{AdS}_5 \times S^5$

- These **localised black holes** have **horizon topology**  $S^8$  and asymptote to  $\text{AdS}_5 \times S^5$ , whose **constant time and 'holographic' slices** have topology  $S^3 \times S^5$ .
- Can think of it as **'inverted' black rings**.


Why is it hard to solve for these solutions?

# Localised Black Holes in $\text{AdS}_5 \times S^5$

- These **localised black holes** have **horizon topology**  $S^8$  and asymptote to  $\text{AdS}_5 \times S^5$ , whose **constant time and 'holographic' slices** have topology  $S^3 \times S^5$ .
- Can think of it as **'inverted' black rings**.

Why is it hard to solve for these solutions?

- Need to solve equations on a pentagon!




# Localised Black Holes in $\text{AdS}_5 \times S^5$

- These **localised black holes** have **horizon topology**  $S^8$  and asymptote to  $\text{AdS}_5 \times S^5$ , whose **constant time and 'holographic' slices** have topology  $S^3 \times S^5$ .
- Can think of it as **'inverted' black rings**.

Why is it hard to solve for these solutions?

- Need to solve equations on a pentagon!
  - Horizon
  - $S^3$  axis
  - South pole
  - North pole




# Localised Black Holes in $\text{AdS}_5 \times S^5$

- These **localised black holes** have **horizon topology**  $S^8$  and asymptote to  $\text{AdS}_5 \times S^5$ , whose **constant time and 'holographic' slices** have topology  $S^3 \times S^5$ .
- Can think of it as **'inverted' black rings**.

Why is it hard to solve for these solutions?


- Need to solve equations on a pentagon!
  - Horizon
  - $S^3$  axis
  - South pole
  - North pole
- They live in 10d - strong decay.






# Lumpy holes in $\text{AdS}_5 \times S^5$ - '15 Dias, Santos and Way

- **Localised holes** in  $\text{AdS}_5 \times S^5$  have a **higher entropy** than smeared Schwarzschild- $\text{AdS}_5$  at **fixed energy**.



Lumpy holes in  $\text{AdS}_5 \times S^5$  - '15 Dias, Santos and Way

- **Localised holes** in  $\text{AdS}_5 \times S^5$  have a **higher entropy** than smeared Schwarzschild- $\text{AdS}_5$  at **fixed energy**.



- They can be the **endpoint of the instability** - modulo violation of weak cosmic censorship!

## Conclusions:

- We have constructed localised black holes in  $\text{AdS}_5 \times S^5$ .
- They are likely to be the endpoint of the GL instability.
- New phase dominates microcanonical ensemble.

## Conclusions:

- We have constructed localised black holes in  $\text{AdS}_5 \times S^5$ .
- They are likely to be the endpoint of the GL instability.
- New phase dominates microcanonical ensemble.

## What should you ask me now:

- Non-uniqueness for smeared Schwarzschild- $\text{AdS}_5$ ?
- What is the story in the canonical ensemble?
- Can you make predictions for field theory observables?

## Conclusions:

- We have constructed localised black holes in  $\text{AdS}_5 \times S^5$ .
- They are likely to be the endpoint of the GL instability.
- New phase dominates microcanonical ensemble.

## What should you ask me now:

- Non-uniqueness for smeared Schwarzschild- $\text{AdS}_5$ ?
- What is the story in the canonical ensemble?
- Can you make predictions for field theory observables?

## Outlook:

- What is the field theory interpretation of this phenomenon?
- Can we run Monte-Carlo simulations of  $\mathcal{N} = 4$  SYM in the microcanonical ensemble and confirm phase diagram?
- ...

Thank You!