

Electromagnetic fields with vanishing scalar invariants¹

Marcello Ortaggio

Institute of Mathematics
Academy of Sciences of the Czech Republic

New York – July 13th, 2016

¹Joint work with V. Pravda, Class. Quantum Grav. 33 (2016) 115010

Contents

- 1 Null electromagnetic fields
- 2 Null gravitational fields
- 3 Main result: VSI p -forms
- 4 Universal solutions

Null electromagnetic fields

Definition [Synge (Silberstein, Bateman, Rainich, Ruse, . . .)]

- ① $F_{ab}F^{ab} = 0 \quad (\Leftrightarrow E^2 - B^2 = 0)$
- ② $F_{ab}{}^*F^{ab} = 0 \quad (\Leftrightarrow \vec{E} \cdot \vec{B} = 0)$

Null electromagnetic fields

Definition [Synge (Silberstein, Bateman, Rainich, Ruse, . . .)]

- ① $F_{ab}F^{ab} = 0 \quad (\Leftrightarrow E^2 - B^2 = 0)$
- ② $F_{ab}{}^*F^{ab} = 0 \quad (\Leftrightarrow \vec{E} \cdot \vec{B} = 0)$

- intrinsically Lorentzian property

Null electromagnetic fields

Definition [Synge (Silberstein, Bateman, Rainich, Ruse, . . .)]

- ① $F_{ab}F^{ab} = 0 \quad (\Leftrightarrow E^2 - B^2 = 0)$
- ② $F_{ab}{}^*F^{ab} = 0 \quad (\Leftrightarrow \vec{E} \cdot \vec{B} = 0)$

- intrinsically Lorentzian property
- plane waves

Null electromagnetic fields

Definition [Synge (Silberstein, Bateman, Rainich, Ruse, . . .)]

- ① $F_{ab}F^{ab} = 0 \quad (\Leftrightarrow E^2 - B^2 = 0)$
- ② $F_{ab}{}^*F^{ab} = 0 \quad (\Leftrightarrow \vec{E} \cdot \vec{B} = 0)$

- intrinsically Lorentzian property
- plane waves
- radiating systems: asymptotically $\mathbf{F} = \frac{\mathbf{N}}{r} + \dots$

Null electromagnetic fields

Definition [Synge (Silberstein, Bateman, Rainich, Ruse, . . .)]

- ① $F_{ab}F^{ab} = 0 \quad (\Leftrightarrow E^2 - B^2 = 0)$
- ② $F_{ab}{}^*F^{ab} = 0 \quad (\Leftrightarrow \vec{E} \cdot \vec{B} = 0)$

- intrinsically Lorentzian property
- plane waves
- radiating systems: asymptotically $\mathbf{F} = \frac{\mathbf{N}}{r} + \dots$
- “ ϵ -property”

Null electromagnetic fields

Definition [Synge (Silberstein, Bateman, Rainich, Ruse, . . .)]

- ① $F_{ab}F^{ab} = 0 \quad (\Leftrightarrow E^2 - B^2 = 0)$
- ② $F_{ab}{}^*F^{ab} = 0 \quad (\Leftrightarrow \vec{E} \cdot \vec{B} = 0)$

- intrinsically Lorentzian property
- plane waves
- radiating systems: asymptotically $\mathbf{F} = \frac{\mathbf{N}}{r} + \dots$
- “ ϵ -property”
- solutions of Maxwell \Rightarrow also solve Born-Infeld's and any NLE!
[Schrödinger'35,'43]

Null gravitational fields

All Riemann scalar invariants $I(\text{Riem})$ vanish:

$$R = 0, \quad R_{ab}R^{ab} = 0, \quad R_{abcd}R^{abcd} = 0, \dots$$

Null gravitational fields

All Riemann scalar invariants $I(\text{Riem})$ vanish:

$$R = 0, \quad R_{ab}R^{ab} = 0, \quad R_{abcd}R^{abcd} = 0, \dots$$

- equivalent to “Riemann type” III (N, O)

Null gravitational fields

All Riemann scalar invariants $I(\text{Riem})$ vanish:

$$R = 0, \quad R_{ab}R^{ab} = 0, \quad R_{abcd}R^{abcd} = 0, \dots$$

- equivalent to “Riemann type” III (N, O)
- in HD: Einstein+type N \Rightarrow also solve

quadratic (\supset Gauss-Bonnet): $G_{ab} + \Lambda_0 g_{ab} + (\text{Riem})^2 + \nabla^2 \text{Ric} = 0$

Lovelock gravity: $G_{ab} + \Lambda_0 g_{ab} + \sum_{k=2} (\text{Riem})^k = 0$

[Pravdová-Pravda'08, Málek-Pravda'11, Reall-Tanahashi-Way'14]

Null gravitational fields

All Riemann scalar invariants $I(\text{Riem})$ vanish:

$$R = 0, \quad R_{ab}R^{ab} = 0, \quad R_{abcd}R^{abcd} = 0, \dots$$

- equivalent to “Riemann type” III (N, O)
- in HD: Einstein+type N \Rightarrow also solve

quadratic (\supset Gauss-Bonnet): $G_{ab} + \Lambda_0 g_{ab} + (\text{Riem})^2 + \nabla^2 \text{Ric} = 0$

Lovelock gravity: $G_{ab} + \Lambda_0 g_{ab} + \sum_{k=2} (\text{Riem})^k = 0$

[Pravdová-Pravda'08, Málek-Pravda'11, Reall-Tanahashi-Way'14]

- certain type N *pp*-waves: solve *any* $\mathcal{L}(\text{Riem}, \nabla \text{Riem}, \dots)$
[Güven'87, Amati-Klimčík'89, Horowitz-Steif'90]

Null gravitational fields

All Riemann scalar invariants $I(\text{Riem})$ vanish:

$$R = 0, \quad R_{ab}R^{ab} = 0, \quad R_{abcd}R^{abcd} = 0, \dots$$

- equivalent to “Riemann type” III (N, O)
- in HD: Einstein+type N \Rightarrow also solve

quadratic (\supset Gauss-Bonnet): $G_{ab} + \Lambda_0 g_{ab} + (\text{Riem})^2 + \nabla^2 \text{Ric} = 0$

Lovelock gravity: $G_{ab} + \Lambda_0 g_{ab} + \sum_{k=2} (\text{Riem})^k = 0$

[Pravdová-Pravda'08, Málek-Pravda'11, Reall-Tanahashi-Way'14]

- certain type N *pp*-waves: solve *any* $\mathcal{L}(\text{Riem}, \nabla \text{Riem}, \dots)$

[Güven'87, Amati-Klimčík'89, Horowitz-Steif'90]

these are in fact **VSI** spacetime: any $I(\text{Riem}, \nabla \text{Riem}, \dots) = 0!$

[Pravda-Pravdová-Coley-Milson'02, Coley-Milson-Pravda-Pravdová'04]

- ① what about VSI Maxwell fields?

$$I(F, \nabla F, \dots) = 0$$

- ① what about VSI Maxwell fields?

$$I(F, \nabla F, \dots) = 0$$

- ② also solutions of higher-derivative NLE?

$$\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$$

old idea [Bopp'40, Podolsky'42]

effective theories motivated by string theory

e.g. [Andreev-Tseytlin'88, Thorlacius'98, Chemissany-Kallosh-Ortin'12, ...]

- ① what about VSI Maxwell fields?

$$I(F, \nabla F, \dots) = 0$$

- ② also solutions of higher-derivative NLE?

$$\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$$

old idea [Bopp'40, Podolsky'42]

effective theories motivated by string theory

e.g. [Andreev-Tseytlin'88, Thorlacius'98, Chemissany-Kallosh-Ortin'12, ...]

We studied this in arbitrary dimension n for a p -form F .

Main result: VSI p -forms

Theorem ([M.O.-Pravda'15])

The following two conditions are equivalent:

- ① a p -form field \mathbf{F} is **VSI** in a spacetime g_{ab}
- ② (a) $\mathbf{F} = \ell \wedge \boldsymbol{\omega}$, with $\ell_a \ell^a = 0 = \ell^{a_1} \omega_{a_1 \dots a_{p-1}}$ (i.e., VSI_0)
(b) $\mathcal{L}_\ell \mathbf{F} = 0$
(c) g_{ab} is “degenerate Kundt”.

Main result: VSI p -forms

Theorem ([M.O.-Pravda'15])

The following two conditions are equivalent:

- ① a p -form field \mathbf{F} is **VSI** in a spacetime g_{ab}
- ②
 - (a) $\mathbf{F} = \ell \wedge \omega$, with $\ell_a \ell^a = 0 = \ell^{a_1} \omega_{a_1 \dots a_{p-1}}$ (i.e., VSI_0)
 - (b) $\mathcal{L}_\ell \mathbf{F} = 0$
 - (c) g_{ab} is “degenerate Kundt”.
- Maxwell equations *not* employed
however, $d\mathbf{F} = 0$ with (2a) \Rightarrow (2b)

Main result: VSI p -forms

Theorem ([M.O.-Pravda'15])

The following two conditions are equivalent:

- ① a p -form field \mathbf{F} is **VSI** in a spacetime g_{ab}
- ② (a) $\mathbf{F} = \ell \wedge \boldsymbol{\omega}$, with $\ell_a \ell^a = 0 = \ell^{a_1} \omega_{a_1 \dots a_{p-1}}$ (i.e., VSI_0)
 (b) $\mathcal{L}_\ell \mathbf{F} = 0$
 (c) g_{ab} is “degenerate Kundt”.

- Maxwell equations *not* employed
 however, $d\mathbf{F} = 0$ with (2a) \Rightarrow (2b)
 - (2c) $\Leftrightarrow \ell$ is geodesic, shearfree, twistfree, expansionfree and
 all $\nabla^k Riem$ are “multiply aligned” with ℓ
 (includes all Einstein-Kundt, Minkowski, (A)dS, . . .)
- [Coley-Hervik-Pelavas'09, Coley-Hervik-Papadopoulos-Pelavas'09]

Main result: VSI p -forms

Theorem ([M.O.-Pravda'15])

The following two conditions are equivalent:

- ① a p -form field \mathbf{F} is **VSI** in a spacetime g_{ab}
- ② (a) $\mathbf{F} = \ell \wedge \boldsymbol{\omega}$, with $\ell_a \ell^a = 0 = \ell^{a_1} \omega_{a_1 \dots a_{p-1}}$ (i.e., VSI_0)
 (b) $\mathcal{L}_\ell \mathbf{F} = 0$
 (c) g_{ab} is “degenerate Kundt”.

- Maxwell equations *not* employed
 however, $d\mathbf{F} = 0$ with (2a) \Rightarrow (2b)
- (2c) $\Leftrightarrow \ell$ is geodesic, shearfree, twistfree, expansionfree and
 all $\nabla^k Riem$ are “multiply aligned” with ℓ
 (includes all Einstein-Kundt, Minkowski, (A)dS, ...)
 [Coley-Hervik-Pelavas'09, Coley-Hervik-Papadopoulos-Pelavas'09]
- proof based on “algebraic VSI theorem” [Hervik'11] and
 boost-weight classification [Milson-Coley-Pravda-Pravdová'05]

Adapted coordinates ($\ell = \partial_r$, $\ell_a dx^a = du$)

$$ds^2 = 2du[dr + H(u, r, x)du + W_\alpha(u, r, x)dx^\alpha] + g_{\alpha\beta}(u, x)dx^\alpha dx^\beta$$
$$F = \frac{1}{(p-1)!} f_{\alpha_1 \dots \alpha_{p-1}}(u, x) du \wedge dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_{p-1}}$$

Adapted coordinates ($\ell = \partial_r$, $\ell_a dx^a = du$)

$$ds^2 = 2du[dr + H(u, r, x)du + W_\alpha(u, r, x)dx^\alpha] + g_{\alpha\beta}(u, x)dx^\alpha dx^\beta$$

$$F = \frac{1}{(p-1)!} f_{\alpha_1 \dots \alpha_{p-1}}(u, x) du \wedge dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_{p-1}}$$

with

$$W_\alpha(u, r, x) = \textcolor{red}{r} W_\alpha^{(1)}(u, x) + W_\alpha^{(0)}(u, x),$$

$$H(u, r, x) = \textcolor{red}{r}^2 H^{(2)}(u, x) + \textcolor{red}{r} H^{(1)}(u, x) + H^{(0)}(u, x).$$

Adapted coordinates ($\ell = \partial_r$, $\ell_a dx^a = du$)

$$ds^2 = 2du[dr + H(u, r, x)du + W_\alpha(u, r, x)dx^\alpha] + g_{\alpha\beta}(u, x)dx^\alpha dx^\beta$$

$$F = \frac{1}{(p-1)!} f_{\alpha_1 \dots \alpha_{p-1}}(u, x) du \wedge dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_{p-1}}$$

with

$$W_\alpha(u, r, x) = \textcolor{red}{r} W_\alpha^{(1)}(u, x) + W_\alpha^{(0)}(u, x),$$

$$H(u, r, x) = \textcolor{red}{r}^2 H^{(2)}(u, x) + \textcolor{red}{r} H^{(1)}(u, x) + H^{(0)}(u, x).$$

- Maxwell equations: f harmonic in $g_{\alpha\beta}$

Adapted coordinates ($\ell = \partial_r$, $\ell_a dx^a = du$)

$$ds^2 = 2du[dr + H(u, r, x)du + W_\alpha(u, r, x)dx^\alpha] + g_{\alpha\beta}(u, x)dx^\alpha dx^\beta$$

$$\mathbf{F} = \frac{1}{(p-1)!} f_{\alpha_1 \dots \alpha_{p-1}}(u, x) du \wedge dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_{p-1}}$$

with $W_\alpha(u, r, x) = \textcolor{red}{r} W_\alpha^{(1)}(u, x) + W_\alpha^{(0)}(u, x)$,

$$H(u, r, x) = \textcolor{red}{r}^2 H^{(2)}(u, x) + \textcolor{red}{r} H^{(1)}(u, x) + H^{(0)}(u, x).$$

- Maxwell equations: f harmonic in $g_{\alpha\beta}$
- Chern-Simons term $\mathbf{F} \wedge \mathbf{F} \wedge \dots$ vanishes identically
(except when linear $n = 2p - 1$)

Adapted coordinates ($\ell = \partial_r$, $\ell_a dx^a = du$)

$$ds^2 = 2du[dr + H(u, r, x)du + W_\alpha(u, r, x)dx^\alpha] + g_{\alpha\beta}(u, x)dx^\alpha dx^\beta$$

$$\mathbf{F} = \frac{1}{(p-1)!} f_{\alpha_1 \dots \alpha_{p-1}}(u, x) du \wedge dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_{p-1}}$$

with $W_\alpha(u, r, x) = \textcolor{red}{r} W_\alpha^{(1)}(u, x) + W_\alpha^{(0)}(u, x)$,

$$H(u, r, x) = \textcolor{red}{r}^2 H^{(2)}(u, x) + \textcolor{red}{r} H^{(1)}(u, x) + H^{(0)}(u, x).$$

- Maxwell equations: \mathbf{f} harmonic in $g_{\alpha\beta}$
- Chern-Simons term $\mathbf{F} \wedge \mathbf{F} \wedge \dots$ vanishes identically
(except when linear $n = 2p - 1$)
- constraints on $W_\alpha^{(1)}$, $W_\alpha^{(0)}$, $H^{(2)}$, $H^{(1)}$, $H^{(0)}$, $g_{\alpha\beta}$ if
backreaction included [Podolský-Žofka'09]

Adapted coordinates ($\ell = \partial_r$, $\ell_a dx^a = du$)

$$ds^2 = 2du[dr + H(u, r, x)du + W_\alpha(u, r, x)dx^\alpha] + g_{\alpha\beta}(u, x)dx^\alpha dx^\beta$$

$$\mathbf{F} = \frac{1}{(p-1)!} f_{\alpha_1 \dots \alpha_{p-1}}(u, x) du \wedge dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_{p-1}}$$

with $W_\alpha(u, r, x) = \textcolor{red}{r} W_\alpha^{(1)}(u, x) + W_\alpha^{(0)}(u, x)$,

$$H(u, r, x) = \textcolor{red}{r}^2 H^{(2)}(u, x) + \textcolor{red}{r} H^{(1)}(u, x) + H^{(0)}(u, x).$$

- Maxwell equations: \mathbf{f} harmonic in $g_{\alpha\beta}$
- Chern-Simons term $\mathbf{F} \wedge \mathbf{F} \wedge \dots$ vanishes identically
(except when linear $n = 2p - 1$)
- constraints on $W_\alpha^{(1)}$, $W_\alpha^{(0)}$, $H^{(2)}$, $H^{(1)}$, $H^{(0)}$, $g_{\alpha\beta}$ if backreaction included [Podolský-Žofka'09]
- waves in Minkowski, (A)dS, Nariai, gyratons, ...

Examples of VSI tensors:

²[M.O.-Pravda'15]

³[Pravda-Pravdová-Coley-Milson'02, Coley-Milson-Pravda-Pravdová'04,
Pelavas-Coley-Milson-Pravda-Pravdová'05]

Examples of VSI tensors:

	vector ℓ	p -form \mathbf{F} ²	Riem ³
VSI ₀	null	N	III (N,O)
VSI ₁	Kundt	N, $\mathcal{L}_\ell \mathbf{F} = 0$, Kundt	N, $\kappa = 0$, $\sigma \Psi_4 = \rho \Phi_{22}$
VSI ₂	Kundt, Riem II	N, $\mathcal{L}_\ell \mathbf{F} = 0$, Kundt, Riem II	III, Kundt
VSI ₃	degKundt	N, $\mathcal{L}_\ell \mathbf{F} = 0$, degKundt	"
:	"	"	"
VSI	"	"	"

²[M.O.-Pravda'15]³[Pravda-Pravdová-Coley-Milson'02, Coley-Milson-Pravda-Pravdová'04, Pelavas-Coley-Milson-Pravda-Pravdová'05]

Universal solutions

Can some VSI F solve any theory $\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$?

Universal solutions

Can some VSI F solve any theory $\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$?

- arbitrary gravity theories $\mathcal{L} = \mathcal{L}(\text{Riem}, \nabla \text{Riem}, \dots)$: solved by “universal” (Einstein) metrics [Coley-Gibbons-Hervik-Pope'08, Hervik-Pravda-Pravdová'14, Hervik-Málek-Pravda-Pravdová'15]

Universal solutions

Can some VSI F solve any theory $\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$?

- arbitrary gravity theories $\mathcal{L} = \mathcal{L}(\text{Riem}, \nabla \text{Riem}, \dots)$: solved by “universal” (Einstein) metrics [Coley-Gibbons-Hervik-Pope'08, Hervik-Pravda-Pravdová'14, Hervik-Málek-Pravda-Pravdová'15]
- Born-Infeld NLE: solved by any null Maxwell field

$$F_{[ab,c]} = 0, \quad \left(\frac{F^{ab} - \mathcal{G}^* F^{ab}}{\sqrt{1 + \mathcal{F} - \mathcal{G}^2}} \right)_{;b} = 0$$

$$(\mathcal{F} \equiv \frac{1}{2} F_{ab} F^{ab}, \mathcal{G} \equiv \frac{1}{4} F_{ab} {}^* F^{ab}) \text{ [Schrödinger'35,'43]}$$

Universal solutions

Can some VSI \mathbf{F} solve any theory $\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$?

- arbitrary gravity theories $\mathcal{L} = \mathcal{L}(\text{Riem}, \nabla \text{Riem}, \dots)$: solved by “universal” (Einstein) metrics [Coley-Gibbons-Hervik-Pope'08, Hervik-Pravda-Pravdová'14, Hervik-Málek-Pravda-Pravdová'15]
- Born-Infeld NLE: solved by any null Maxwell field

$$F_{[ab,c]} = 0, \quad \left(\frac{F^{ab} - \mathcal{G}^* F^{ab}}{\sqrt{1 + \mathcal{F} - \mathcal{G}^2}} \right)_{;b} = 0$$

$$(\mathcal{F} \equiv \frac{1}{2} F_{ab} F^{ab}, \mathcal{G} \equiv \frac{1}{4} F_{ab} {}^* F^{ab}) \text{ [Schrödinger'35,'43]}$$

- arbitrary $\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$: any VSI \mathbf{F} in type III Kundt

Universal solutions

Can some VSI \mathbf{F} solve any theory $\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$?

- arbitrary gravity theories $\mathcal{L} = \mathcal{L}(\text{Riem}, \nabla \text{Riem}, \dots)$: solved by “universal” (Einstein) metrics [Coley-Gibbons-Hervik-Pope'08, Hervik-Pravda-Pravdová'14, Hervik-Málek-Pravda-Pravdová'15]
- Born-Infeld NLE: solved by any null Maxwell field

$$F_{[ab,c]} = 0, \quad \left(\frac{F^{ab} - \mathcal{G}^* F^{ab}}{\sqrt{1 + \mathcal{F} - \mathcal{G}^2}} \right)_{;b} = 0$$

$$(\mathcal{F} \equiv \frac{1}{2} F_{ab} F^{ab}, \mathcal{G} \equiv \frac{1}{4} F_{ab} {}^* F^{ab}) \quad [\text{Schrödinger}'35, '43]$$

- arbitrary $\mathcal{L} = \mathcal{L}(F, \nabla F, \dots)$: any VSI \mathbf{F} in type III Kundt
- coupling to gravity also possible: further restrictions
cf. also [Güven'87, Horowitz-Steif'90, Coley'02]
NLE in [Kichenassamy'59, Kremer-Kichenassamy'60, Peres'60]

An Einstein-Maxwell universal example ($n = 4, p = 2$):

$$\begin{aligned}ds^2 &= 2du[dr + \frac{1}{2}(xr - xe^x - 2\kappa_0 e^x c^2(u))du] + e^x(dx^2 + e^{2u}dy^2) \\F &= e^{x/2}c(u)du \wedge \left(-\cos \frac{ye^u}{2}dx + e^u \sin \frac{ye^u}{2}dy\right)\end{aligned}$$

- $\ell = \partial_r$ is Kundt and *recurrent* ($\ell_{a;b} = f\ell_a\ell_b$)
- F is VSI
- Petrov type III
- obtained by “charging” a vacuum metric of Petrov [Petrov'62]