



**21st International Conference
on General Relativity
and Gravitation**
Columbia University, New York



Spontaneous scalarization of neutron stars in the unconstrained parameter regime of scalar-tensor theories

(Based on arXiv:1604.04175)

Néstor Ortiz

In collaboration with Raissa Mendes

**Perimeter Institute for Theoretical Physics
Waterloo, ON, Canada.**

GR21. New York City. July 11th, 2016.

Motivation

From Raissa's talk we have learned:

In Scalar-Tensor Theories (STT) with $\beta_0 > 0$, highly compact neutron stars (NS) obeying realistic equations of state (EoS) possess linearly unstable scalar modes.

Motivation

From Raissa's talk we have learned:

In Scalar-Tensor Theories (STT) with $\beta_0 > 0$, highly compact neutron stars (NS) obeying realistic equations of state (EoS) possess linearly unstable scalar modes.

Brief review

In the Einstein frame: $S = \frac{1}{16\pi} \int d^4x \sqrt{-g} (R - 2\nabla_\mu \phi \nabla^\mu \phi) + S_m[\Psi_m; a(\phi)^2 g_{\mu\nu}]$

Coupling function 

Motivation

From Raissa's talk we have learned:

In Scalar-Tensor Theories (STT) with $\beta_0 > 0$, highly compact neutron stars (NS) obeying realistic equations of state (EoS) possess linearly unstable scalar modes.

Brief review

In the Einstein frame: $S = \frac{1}{16\pi} \int d^4x \sqrt{-g} (R - 2\nabla_\mu \phi \nabla^\mu \phi) + S_m[\Psi_m; a(\phi)^2 g_{\mu\nu}]$

$$\beta_0 := \left. \frac{d\alpha}{d\phi} \right|_{\phi_0} \quad \leftarrow \quad \alpha(\phi) := \frac{d}{d\phi} \ln a(\phi) \quad \leftarrow \text{Coupling function} \quad \nearrow$$

Motivation

From Raissa's talk we have learned:

In Scalar-Tensor Theories (STT) with $\beta_0 > 0$, highly compact neutron stars (NS) obeying realistic equations of state (EoS) possess linearly unstable scalar modes.

Brief review

In the Einstein frame: $S = \frac{1}{16\pi} \int d^4x \sqrt{-g} (R - 2\nabla_\mu \phi \nabla^\mu \phi) + S_m[\Psi_m; a(\phi)^2 g_{\mu\nu}]$

$$\beta_0 := \left. \frac{d\alpha}{d\phi} \right|_{\phi_0} \quad \leftarrow \quad \alpha(\phi) := \frac{d}{d\phi} \ln a(\phi) \quad \leftarrow \text{Coupling function} \quad \nearrow$$

NS: spherically symmetric perfect fluids, i. e.

$$T^{\mu\nu} = \epsilon u^\mu u^\nu + p(\tilde{g}^{\mu\nu} + u^\mu u^\nu)$$

By highly compact we mean $[p > \epsilon/3]_{star \text{ center}}$

Motivation

From Raissa's talk we have learned:

In Scalar-Tensor Theories (STT) with $\beta_0 > 0$, highly compact neutron stars (NS) obeying realistic equations of state (EoS) possess linearly unstable scalar modes.

Brief review


In the Einstein frame: $S = \frac{1}{16\pi} \int d^4x \sqrt{-g} (R - 2\nabla_\mu \phi \nabla^\mu \phi) + S_m[\Psi_m; a(\phi)^2 g_{\mu\nu}]$

$$\beta_0 := \left. \frac{d\alpha}{d\phi} \right|_{\phi_0} \quad \leftarrow \quad \alpha(\phi) := \frac{d}{d\phi} \ln a(\phi)$$

NS: spherically symmetric perfect fluids, i. e.

$$T^{\mu\nu} = \epsilon u^\mu u^\nu + p(\tilde{g}^{\mu\nu} + u^\mu u^\nu)$$

By highly compact we mean $[p > \epsilon/3]_{star\ center}$



Motivation

From Raissa's talk we have learned:

In Scalar-Tensor Theories (STT) with $\beta_0 > 0$, highly compact neutron stars (NS) obeying realistic equations of state (EoS) possess linearly unstable scalar modes.

Our goal

To investigate the final fate of such unstable equilibrium stars

Do they evolve to a stable, scalarized state? Do they collapse?

Motivation

From Raissa's talk we have learned:

In Scalar-Tensor Theories (STT) with $\beta_0 > 0$, highly compact neutron stars (NS) obeying realistic equations of state (EoS) possess linearly unstable scalar modes.

Our goal

To investigate the final fate of such unstable equilibrium stars

Do they evolve to a stable, scalarized state? Do they collapse?

Our strategies

- * Energy balance analysis
- * Full non-linear numerical simulations

Posing problem...

What we want to solve

Field equations

Fluid equations

$$\left\{ \begin{array}{l} G_{\mu\nu} - 2\nabla_\mu\phi\nabla_\nu\phi + g_{\mu\nu}\nabla_\sigma\phi\nabla^\sigma\phi = 8\pi a^2 T_{\mu\nu} \\ \nabla^\mu\nabla_\mu\phi = -4\pi a^4\alpha T \end{array} \right.$$
$$\left\{ \begin{array}{l} \nabla_\mu(\rho u^\mu) = 0 \\ \nabla_\nu T^{\mu\nu} = 0 \end{array} \right.$$

Posing problem...

What we want to solve

Field equations

Fluid equations

$$\left\{ \begin{array}{l} G_{\mu\nu} - 2\nabla_\mu\phi\nabla_\nu\phi + g_{\mu\nu}\nabla_\sigma\phi\nabla^\sigma\phi = 8\pi a^2 T_{\mu\nu} \\ \nabla^\mu\nabla_\mu\phi = -4\pi a^4\alpha T \end{array} \right.$$

$$\left\{ \begin{array}{l} \nabla_\mu(\rho u^\mu) = 0 \\ \nabla_\nu T^{\mu\nu} = 0 \end{array} \right.$$

+ Polytropic EoS

$$\begin{aligned} p(\rho) &= K\rho_0(\rho/\rho_0)^\gamma \\ \gamma &= 3, K = 0.005, \\ \rho_0 &= 1.66 \times 10^{14} \text{g/cm}^3 \end{aligned}$$

Posing problem...

What we want to solve

Field equations

Fluid equations

$$\begin{cases} G_{\mu\nu} - 2\nabla_\mu\phi\nabla_\nu\phi + g_{\mu\nu}\nabla_\sigma\phi\nabla^\sigma\phi = 8\pi a^2 T_{\mu\nu} \\ \nabla^\mu\nabla_\mu\phi = -4\pi a^4\alpha T \end{cases}$$
$$\begin{cases} \nabla_\mu(\rho u^\mu) = 0 \\ \nabla_\nu T^{\mu\nu} = 0 \end{cases}$$

+ Polytropic EoS

$$p(\rho) = K\rho_0(\rho/\rho_0)^\gamma$$
$$\gamma = 3, K = 0.005,$$
$$\rho_0 = 1.66 \times 10^{14} \text{g/cm}^3$$

Our approach: **3 + 1 Formalism, spherical symmetry** (reduction to 1+1D problem)

Radial gauge / Polar slicing $ds^2 = -N(t, r)^2 dt^2 + A(t, r)^2 dr^2 + r^2(d\vartheta^2 + \sin^2\vartheta d\varphi^2)$

Posing problem...

What we want to solve

Field equations

Fluid equations

$$\begin{cases} G_{\mu\nu} - 2\nabla_\mu\phi\nabla_\nu\phi + g_{\mu\nu}\nabla_\sigma\phi\nabla^\sigma\phi = 8\pi a^2 T_{\mu\nu} \\ \nabla^\mu\nabla_\mu\phi = -4\pi a^4\alpha T \end{cases}$$

$$\begin{cases} \nabla_\mu(\rho u^\mu) = 0 \\ \nabla_\nu T^{\mu\nu} = 0 \end{cases}$$

+ Polytropic EoS

$$p(\rho) = K\rho_0(\rho/\rho_0)^\gamma$$

$$\gamma = 3, K = 0.005,$$

$$\rho_0 = 1.66 \times 10^{14} \text{g/cm}^3$$

Our approach: **3 + 1 Formalism, spherical symmetry** (reduction to 1+1D problem)

Radial gauge / Polar slicing $ds^2 = -N(t, r)^2 dt^2 + A(t, r)^2 dr^2 + r^2(d\vartheta^2 + \sin^2\vartheta d\varphi^2)$

Numerical methods: **Finite volume + High Resolution Shock-Capturing scheme**

Requires fluid equations in flux-conservative form: $\frac{\partial}{\partial t}(A\mathbf{q}) + \frac{1}{r^2} \frac{\partial}{\partial r} (N A r^2 \mathbf{F}(\mathbf{q})) = \mathbf{S}(\mathbf{q})$

Posing problem...

What we want to solve

Field equations

$$\begin{cases} G_{\mu\nu} - 2\nabla_\mu\phi\nabla_\nu\phi + g_{\mu\nu}\nabla_\sigma\phi\nabla^\sigma\phi = 8\pi a^2 T_{\mu\nu} \\ \nabla^\mu\nabla_\mu\phi = -4\pi a^4\alpha T \end{cases}$$

Fluid equations

$$\begin{cases} \nabla_\mu(\rho u^\mu) = 0 \\ \nabla_\nu T^{\mu\nu} = 0 \end{cases}$$

+ Polytropic EoS

$$\begin{aligned} p(\rho) &= K\rho_0(\rho/\rho_0)^\gamma \\ \gamma &= 3, K = 0.005, \\ \rho_0 &= 1.66 \times 10^{14} \text{g/cm}^3 \end{aligned}$$

Our approach: **3 + 1 Formalism, spherical symmetry** (reduction to 1+1D problem)

Radial gauge / Polar slicing $ds^2 = -N(t, r)^2 dt^2 + A(t, r)^2 dr^2 + r^2(d\vartheta^2 + \sin^2\vartheta d\varphi^2)$

Numerical methods: **Finite volume + High Resolution Shock-Capturing scheme**

Requires fluid equations in flux-conservative form: $\frac{\partial}{\partial t}(A\mathbf{q}) + \frac{1}{r^2}\frac{\partial}{\partial r}(NAr^2\mathbf{F}(\mathbf{q})) = \mathbf{S}(\mathbf{q})$

Coupling function:

Model 1 (M1): $\alpha(\phi) = \tanh[\sqrt{3}\beta(\phi - \phi_0)]/\sqrt{3}$
[mimics Nonminimally coupled (NMC) field]


Model 2 (M2): $\alpha(\phi) = \beta(\phi - \phi_0)$
[Damour-Esposito-Farese (DEF)]

Relevant previous results

(Dynamic transition to scalarized states in the $\beta < 0$ case)

Nonminimally coupled fields (like M1): M. Alcubierre et al., 2010, M. Ruiz, 2012.

Damour-Esposito-Farese case (M2), J. Novak, 1998



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $	$ \omega [M_\odot]$
1*	10.0	2.3594	1.9650	0.287	0	0
2 (M1)	9.2061	2.3594	1.9641	0.273	0.095	0.30
3 (M2)	8.0747	2.3594	1.9459	0.209	0.284	1.23


Table 1: Some equilibrium solutions with $\beta = -6$. The solution marked with a star is used as initial data for our numerical simulation.

Relevant previous results

(Dynamic transition to scalarized states in the $\beta < 0$ case)

Nonminimally coupled fields (like M1): M. Alcubierre et al., 2010, M. Ruiz, 2012.

Damour-Esposito-Farese case (M2), J. Novak, 1998



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $	$ \omega [M_\odot]$
1*	10.0	2.3594	1.9650	0.287	0	0
2 (M1)	9.2061	2.3594	1.9641	0.273	0.095	0.30
3 (M2)	8.0747	2.3594	1.9459	0.209	0.284	1.23

Table 1: Some equilibrium solutions with $\beta = -6$. The solution marked with a star is used as initial data for our numerical simulation.


Solutions 2 and 3 are energetically favored!

Relevant previous results

(Dynamic transition to scalarized states in the $\beta < 0$ case)

Nonminimally coupled fields (like M1): M. Alcubierre et al., 2010, M. Ruiz, 2012.


Damour-Esposito-Farese case (M2), J. Novak, 1998



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $	$ \omega [M_\odot]$
1*	10.0	2.3594	1.9650	0.287	0	0
2 (M1)	9.2061	2.3594	1.9641	0.273	0.095	0.30
3 (M2)	8.0747	2.3594	1.9459	0.209	0.284	1.23

Table 1: Some equilibrium solutions with $\beta = -6$. The solution marked with a star is used as initial data for our numerical simulation.

Solutions 2 and 3 are energetically favored!




The transition is consistent with solutions 2 and 3 in the Table 1

Relevant previous results

(Dynamic transition to scalarized states in the $\beta < 0$ case)

Nonminimally coupled fields (like M1): M. Alcubierre et al., 2010, M. Ruiz, 2012.


Damour-Esposito-Farese case (M2), J. Novak, 1998



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $	$ \omega [M_\odot]$
1*	10.0	2.3594	1.9650	0.287	0	0
2 (M1)	9.2061	2.3594	1.9641	0.273	0.095	0.30
3 (M2)	8.0747	2.3594	1.9459	0.209	0.284	1.23

Table 1: Some equilibrium solutions with $\beta = -6$. The solution marked with a star is used as initial data for our numerical simulation.

Solutions 2 and 3 are energetically favored!




The transition is consistent with solutions 2 and 3 in the Table 1

Remark: The transition is generic regardless of the details of coupling function

New results

Exploring the $\beta > 0$ case

Model 1
(NMC field)




Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827

Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.

New results

Exploring the $\beta > 0$ case

Model 1
(NMC field)



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827


Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.

Solution 2 is energetically favored!



New results

Exploring the $\beta > 0$ case

Model 1
(NMC field)



Dynamic transition to scalarized state!
(consistent with solution 2 the Table 2)



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827


Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.

Solution 2 is energetically favored!



New results

Exploring the $\beta > 0$ case

Model 1
(NMC field)



Dynamic transition to scalarized state!
(consistent with solution 2 the Table 2)



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827

Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.


Solution 2 is energetically favored!

Spontaneous scalarization can happen in theories with $\beta > 0$

New results

Exploring the $\beta > 0$ case

Model 1
(NMC field)




Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827
5*	12.0	2.4529	2.02461	0.309	0
6	13.4600	2.4529	2.02469	0.320	0.017240

Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.

New results

Exploring the $\beta > 0$ case

Model 1
(NMC field)



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827
5*	12.0	2.4529	2.02461	0.309	0
6	13.4600	2.4529	2.02469	0.320	0.017240


Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.

Solution 6 is energetically disfavored vs. solution 5: likely unphysical


New results

Exploring the $\beta > 0$ case

Model 1
(NMC field)




Gravitational collapse!



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827
5*	12.0	2.4529	2.02461	0.309	0
6	13.4600	2.4529	2.02469	0.320	0.017240

Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.




Solution 6 is energetically disfavored vs. solution 5: likely unphysical


New results

Exploring the $\beta > 0$ case

Model 1
(NMC field)




Gravitational collapse!



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827
5*	12.0	2.4529	2.02461	0.309	0
6	13.4600	2.4529	2.02469	0.320	0.017240

Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.




Unbounded growths in $r = 0$

Solution 6 is energetically disfavored vs. solution 5: likely unphysical

New results

Exploring the $\beta > 0$ case





Model 1
(NMC field)



Gravitational collapse!

Slice stretching

Lapse function collapses



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	11.4251	2.4304	2.01053	0.304	0.023159
3	12.2279	2.4304	2.01056	0.311	0.039036
4	14.3890	2.4304	2.01132	0.325	0.027827
5*	12.0	2.4529	2.02461	0.309	0
6	13.4600	2.4529	2.02469	0.320	0.017240

Table 2: Some equilibrium solutions in M1 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.

Unbounded growths in $r = 0$


Solution 6 is energetically disfavored vs. solution 5: likely unphysical

New results

Exploring the $\beta > 0$ case

Model 2
(DEF)

(In agreement with C. Palenzuela and S. Liebling, 2016)




New results

Exploring the $\beta > 0$ case

Model 2
(DEF)

(In agreement with C. Palenzuela and S. Liebling, 2016)



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	10.9493	2.4304	2.01087	0.312	0.055
3	14.4841	2.4304	2.01142	0.327	0.023
4*	11.20	2.43611	2.01435	0.313	0.050
5*	11.20	2.39877	1.99273	0.325	0.081
6*	11.20	2.32321	1.94928	0.332	0.104


Table 3: Some equilibrium solutions in M2 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.

New results

Exploring the $\beta > 0$ case

Model 2
(DEF)

(In agreement with C. Palenzuela and S. Liebling, 2016)



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	10.9493	2.4304	2.01087	0.312	0.055
3	14.4841	2.4304	2.01142	0.327	0.023
4*	11.20	2.43611	2.01435	0.313	0.050
5*	11.20	2.39877	1.99273	0.325	0.081
6*	11.20	2.32321	1.94928	0.332	0.104

Table 3: Some equilibrium solutions in M2 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.


Solutions 2 and 3 are energetically disfavored vs. solution 1: likely unphysical

New results


Exploring the $\beta > 0$ case

Model 2
(DEF)

(In agreement with C. Palenzuela and S. Liebling, 2016)




Gravitational collapse!



Solution	$\tilde{\rho}_c/\rho_0$	$M_b[M_\odot]$	$M[M_\odot]$	M/R_s	$ \phi_c - \phi_0 $
1*	11.20	2.4304	2.01058	0.302	0
2	10.9493	2.4304	2.01087	0.312	0.055
3	14.4841	2.4304	2.01142	0.327	0.023
4*	11.20	2.43611	2.01435	0.313	0.050
5*	11.20	2.39877	1.99273	0.325	0.081
6*	11.20	2.32321	1.94928	0.332	0.104

Table 3: Some equilibrium solutions in M2 with $\beta = 100$. Solutions marked with stars are used as initial data for numerical simulations.



Solutions 2 and 3 are energetically disfavored vs. solution 1: likely unphysical

Conclusions

Punch line:

First demonstration of spontaneous scalarization in STT with $\beta_0 > 0$

Conclusions

Punch line:

First demonstration of spontaneous scalarization in STT with $\beta_0 > 0$

Requires:



Highly compact NS's, i. e.

$$[p > \epsilon/3]_{star\ center}$$

Conclusions

Punch line:

First demonstration of spontaneous scalarization in STT with $\beta_0 > 0$

Caveat:

Final state is highly sensitive to the coupling function, varying from gravitational collapse to spontaneous scalarization

Requires:

Highly compact NS's, i. e.
 $[p > \epsilon/3]_{star\ center}$

Conclusions

Punch line:

First demonstration of spontaneous scalarization in STT with $\beta_0 > 0$

Caveat:

Final state is highly sensitive to the coupling function, varying from gravitational collapse to spontaneous scalarization

Requires:

Highly compact NS's, i. e.
 $[p > \epsilon/3]_{star\ center}$

Constraints:

Would be possible when NS with $M/R \gtrsim 0.27$ are confirmed

Conclusions

Punch line:

First demonstration of spontaneous scalarization in STT with $\beta_0 > 0$

Caveat:

Final state is highly sensitive to the coupling function, varying from gravitational collapse to spontaneous scalarization

Requires:

Highly compact NS's, i. e.
 $[p > \epsilon/3]_{star\ center}$

Constraints:

Would be possible when NS with $M/R \gtrsim 0.27$ are confirmed

Bounds on β_0 should be attributed to particular models. Thus, observations can potentially be used to probe the structure of the coupling function

Conclusions

Punch line:

First demonstration of spontaneous scalarization in STT with $\beta_0 > 0$

Caveat:

Final state is highly sensitive to the coupling function, varying from gravitational collapse to spontaneous scalarization

Requires:

Highly compact NS's, i. e.
 $[p > \epsilon/3]_{star\ center}$

Constraints:

Would be possible when NS with $M/R \gtrsim 0.27$ are confirmed

Bounds on β_0 should be attributed to particular models. Thus, observations can potentially be used to probe the structure of the coupling function