

Can chaos be observed in quantum gravity?

Philipp Höhn

Institute for Quantum Optics and Quantum Information, Vienna, and
Vienna Center for Quantum Science and Technology

GR 21, New York
11 July 2016

based on:

PH, M. Kubalova and A. Tsobanjan, PRD **86** 065014 (2012);
B. Dittrich, PH, T. Koslowski and M. Nelson arXiv:1508.01947, and
arXiv:1602.03237

GR and 'observables'

General Relativity is a gauge theory

⇒ physical observables should be diffeomorphism invariant

canonically:

- Dirac observables as 'constants of motion' of constraints
- dynamics relationally ⇒ 'evolving constants of motion' [Wheeler 60's; Rovelli 90's; Dittrich '06, '07,.....]

⇒ notoriously difficult to construct

often overlooked: even absent in presence of chaos

- 1 what then is observable?
- 2 consequences for QT?

Non-integrability and constraints [Dittrich, PH, Koslowski, Nelson '15; '16]

recall: non-integrable (unconstrained) systems:

- no global (smooth) constants of motion other than H exist
- ⇒ trajectories lie on $(2N - 1)$ -dim. energy surface

now: constrained system weakly non-integrable if:

- ⚡ differentiable Dirac observables indep. of constraints
- ⇒ ⚡ reduced phase space
- ⇒ gauge invariant DoFs exist, but non-differentiable (or local)
- ⇒ no Poisson algebraic structure ⇒ **how to represent observables in QT?**

difference:

unconstrained: do not need to solve dynamics

constrained: need to solve dynamics

to access physical DoFs

Why worry about this?

GR likely weakly non-integrable (N-body problem, Mixmaster, BKL,...)

- ⇒ (probably) \nexists smooth Dirac observables and reduced phase space in full GR
- ⇒ what are repercussions for QG?

Toy model: free particles on a circle [Dittrich, PH, Koslowski, Nelson '15; '16]

Compactify free dynamics: $x_i + 1 \sim x_i$, $i = 1, 2 \Rightarrow$ conf. manif. $\mathcal{Q} \simeq T^2$

$$C = \frac{p_1^2}{2m_1} + \frac{p_2^2}{2m_2} - E \approx 0$$

■ solutions to EoMs


$$x_i(t) = \frac{p_i}{m_i} t + x_{i0} - n_i$$

$n_i := \lfloor \frac{p_i}{m_i} t + x_{i0} \rfloor$ winding number in x_i

if:

$\frac{m_2}{m_1} \frac{p_1}{p_2} \in \mathbb{Q}$: resonant torus, periodic orbits


$\frac{m_2}{m_1} \frac{p_1}{p_2} \notin \mathbb{Q}$: non-resonant torus, **ergodic** orbits



Absence of sufficiently many Dirac observables [Dittrich, PH, Kosłowski, Nelson '15; '16]

- momenta p_i are Dirac observables
- BUT: $O(p_i; x_1, x_2)$ const. on trajectories with $\partial_i O \neq 0$ discontinuous in x_i

- ergodicity destroys integrability



- space of solutions not a manifold
- ⇒ no reduced phase space, no (sufficient) algebra of observables

- can still have gauge invariant 'observables', however, either
 - 1 global and discontinuous, e.g.

$$M = (x_1 + n_1)p_2/m_2 - (x_2 + n_2)p_1/m_1$$

- 2 local [Bojowald, PH, Tsobanjan '11a; '11b]

- $\mathcal{H}_{\text{kin}} = L^2(T^2)$ basis:



$$\psi_{\vec{k}_1, \vec{k}_2}(x_1, x_2) = \exp(2\pi i k_1 x_1) \exp(2\pi i k_2 x_2), \quad \vec{k} \in \mathbb{Z}^2$$

- $\hat{p}_i \psi = -i\hbar \partial_i \psi$

- quantum constraint

$$k_1^2 + \frac{m_1}{m_2} k_2^2 = \frac{2m_1 E}{\hbar^2}$$

- for $m_1/m_2 \notin \mathbb{Q}$ $0 \leq \dim \mathcal{H}_{\text{phys}} \leq 4$



NOT peaked on class. orbit for $m_1/m_2 \notin \mathbb{Q}$

width/separation ≈ 1

- physical transition amplitudes show no semiclassical behaviour

Polymer type quantization: discrete topology [Dittrich, PH, Kosłowski, Nelson '15; '16]

- \mathcal{H}_{kin} given by (uncountable) basis

$$\psi_{x'_1, x'_2}(x_1, x_2) = \delta_{x'_1, x_1} \delta_{x'_2, x_2}$$

- no momenta, but translations

$$(R_1^\mu \psi)(x_1, x_2) = \psi(x_1 + \mu, x_2), \quad (R_2^\mu \psi)(x_1, x_2) = \psi(x_1, x_2 + \mu)$$

- constraint acquires **continuous** spectrum for $\mu \notin \mathbb{Q}$:

$$\left\{ \frac{\hbar^2}{\mu^2} (2 - \cos(2\pi\rho_1) - \cos(2\pi\rho_2)) - E \mid \rho_1, \rho_2 \in [0, 1) \right\}$$

⇒ upon superselec. get ∞ -dim. separable $\mathcal{H}_{\text{phys}}$ as L^2 over 'momentum' ρ


⇒ on this $\mathcal{H}_{\text{phys}}$ have sufficiently many observables

$$\hat{M} := \frac{i}{2\pi} \left(\sin(2\pi\rho_2) \frac{\partial}{\partial \rho_1} - \sin(2\pi\rho_1) \frac{\partial}{\partial \rho_2} \right) \quad (\text{'angular mom.'})$$

$$[\hat{M}, e^{2\pi i \rho_1}] = -e^{2\pi i \rho_1} \sin(2\pi\rho_2)$$

⇒ features good semiclassical transition amplitudes

Closed FRW with (min. coupled) massive scalar [PH, Kubalova, Tsobanjan, '12]



(a) benign solution, (b) defocussing of nearby trajectories in turning region, (c) devoid of global clocks

- Ham. constraint $C = p_\phi^2 - p_\alpha^2 - e^{4\alpha} + m^2 \phi^2 e^{6\alpha}$
- **model chaotic and non-integrable** [Page '84, Cornish, Shellard '98; Belinsky, Khalatnikov, Grishchuk, Zeldovich '85]
- in region of max. expansion α_{max} (chaotic scattering):
 - 1 breakdown of semiclassicality
[also indep. observed in Kiefer '88]
 - 2 relational observables only transient
 \Rightarrow relational evolution breaks down

Conclusions

- Chaos precludes smooth Dirac observables
- ⇒ probably no smooth Dirac observables and red. phase space for full GR
- serious problem for 'standard' constraint quantization

what do we do?

- always \exists generalized discontinuous 'observables'
- ⇒ adapt method of quantization, refine topology until sufficiently many observables continuous
- ⇒ here: polymer quantization overcomes troubles of 'standard' quantization!

further reading: [arXiv:1602.03237](#), [1508.01947](#), [1111.5193](#)