

# Projective State Spaces for Quantum Gravity

Suzanne Lanéry

Centro de Ciencias Matemáticas  
Universidad Nacional Autónoma de México, Morelia



CENTRO DE CIENCIAS  
MATEMÁTICAS

# Introduction

## Quantum Field Theory without a Vacuum State

Standard approach to QFT:

- ▶ discrete excitations around a **vacuum** state
- ▶  $\neq$  vacuum states  $\leftrightarrow$  **inequivalent** representations  $\Rightarrow$  need to fine-tune the vacuum state to the **dynamics**
  - ↳ QFT on curved space time: no natural choice
  - ↳ LQG: preferred kinematical vacuum [Lewandowski, Okołów, Sahlmann, Thiemann; Fleischhack] but tension with the dynamics and the semi-classical limit [see also: Koslowski & Sahlmann '11]

An alternative way to construct the state space [Kijowski '76, Okołów '09 & '13], by gluing together elementary **building blocs**:

- ▶ **lifting** the Stone-von Neumann theorem to QFT
- ▶ yields **universal** quantum state spaces, independent of any choice of **polarization**

# Contents

## Projective Formalism for QFT

Building Blocs: Finitely many Degrees of Freedom

Coarse-graining between Phase Spaces

Coarse-graining between Quantum State Spaces

Natural Embedding of Arbitrary Vacuum Sectors


## Universal Selections of Degrees of Freedom

## Application to Loop Quantum Gravity

# Projective Formalism for QFT

Building Blocs: Finitely many Degrees of Freedom

$\mathcal{M}$



$$a = \frac{\hat{x} + i\hat{p}}{\sqrt{2}}$$

$$|n\rangle = \frac{a^{\dagger n}}{\sqrt{n!}} |\emptyset\rangle$$

Example: linear system

- modes = **independant** pairs of canonically **conjugate** variables
- polarization  $\leadsto$  Fock representation

Relating different choices:


- finite dim  $\Rightarrow$  unitarily equivalent representations (but  $\neq$  vacuum states)
- infinite dim  $\Rightarrow$  in general, **not** equivalent

[Uniqueness for finite dim linear: Stone, von Neumann; Geometric quantization: Woodhouse,...]

# Projective Formalism for QFT

Building Blocs: Finitely many Degrees of Freedom


$\mathcal{M}$



$$a = \frac{\hat{x} + i\hat{p}}{\sqrt{2}}$$

$$|n\rangle = \frac{a^{\dagger n}}{\sqrt{n!}} |\emptyset\rangle$$

$\mathcal{M}$



Example: linear system

- modes = **independant** pairs of canonically **conjugate** variables
- polarization  $\rightsquigarrow$  Fock representation

Relating different choices:


- finite dim  $\Rightarrow$  unitarily equivalent representations (but  $\neq$  vacuum states)
- infinite dim  $\Rightarrow$  in general, **not** equivalent

[Uniqueness for finite dim linear: Stone, von Neumann; Geometric quantization: Woodhouse,...]

# Projective Formalism for QFT

Building Blocs: Finitely many Degrees of Freedom


$\mathcal{M}$



$$a = \frac{\hat{x} + i\hat{p}}{\sqrt{2}}$$

$$|n\rangle = \frac{a^{\dagger n}}{\sqrt{n!}} |\emptyset\rangle$$

$\mathcal{M}$



$$a' = \alpha a + \beta a^\dagger$$

$$|\emptyset'\rangle = \sum_p \xi_{2p} |2p\rangle$$

$$|\alpha|^2 - |\beta|^2 = 1, \quad \xi_{2p} := \left(1 - (\beta/\alpha)^2\right)^{1/4} \frac{\sqrt{(2p)!}}{p!} \left(-\frac{\beta}{2\alpha}\right)^p$$

Example: linear system

- modes = **independant** pairs of canonically **conjugate** variables
- polarization  $\rightsquigarrow$  Fock representation


Relating different choices:

- finite dim  $\Rightarrow$  unitarily equivalent representations (but  $\neq$  vacuum states)
- infinite dim  $\Rightarrow$  in general, **not** equivalent

[Uniqueness for finite dim linear: Stone, von Neumann; Geometric quantization: Woodhouse,...]

# Projective Formalism for QFT

Coarse-graining between Phase Spaces



$$\eta \preccurlyeq \eta' \in \mathcal{L}$$

Collection of partial theories:

- ▶  $\eta \in \mathcal{L}$  = a selection of d.o.f.'s
- ▶ 'small' **phase** space  $\mathcal{M}_\eta$


Coarse graining:

- ▶ projections  $\pi_{\eta' \rightarrow \eta}$  for  $\eta \preccurlyeq \eta'$
- ▶ lifting observables  
 $A_\eta \circ \pi_{\eta' \rightarrow \eta} =: A_{\eta'}$   
↳ compatible with the Poisson brackets
- ▶ **preferred** factorization  
 $\mathcal{M}_{\eta'} \approx \mathcal{M}_{\eta' \rightarrow \eta} \times \mathcal{M}_\eta$

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective Formalism for QFT

Coarse-graining between Phase Spaces



$$\eta \preccurlyeq \eta' \in \mathcal{L}$$

Collection of partial theories:

- $\eta \in \mathcal{L}$  = a selection of d.o.f.'s
- 'small' **phase** space  $\mathcal{M}_\eta$


Coarse graining:

- projections  $\pi_{\eta' \rightarrow \eta}$  for  $\eta \preccurlyeq \eta'$
- lifting observables  
 $A_\eta \circ \pi_{\eta' \rightarrow \eta} =: A_{\eta'}$   
↳ compatible with the Poisson brackets
- **preferred** factorization  
 $\mathcal{M}_{\eta'} \approx \mathcal{M}_{\eta' \rightarrow \eta} \times \mathcal{M}_\eta$

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective Formalism for QFT

Coarse-graining between Phase Spaces



Collection of partial theories:

- $\eta \in \mathcal{L}$  = a selection of d.o.f.'s
- 'small' **phase** space  $\mathcal{M}_\eta$


Coarse graining:

- projections  $\pi_{\eta' \rightarrow \eta}$  for  $\eta \preccurlyeq \eta'$
- lifting observables  
 $A_\eta \circ \pi_{\eta' \rightarrow \eta} =: A_{\eta'}$   
↳ compatible with the Poisson brackets
- **preferred** factorization  
 $\mathcal{M}_{\eta'} \approx \mathcal{M}_{\eta' \rightarrow \eta} \times \mathcal{M}_\eta$

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective Formalism for QFT

Coarse-graining between Phase Spaces



Collection of partial theories:

- ▶  $\eta \in \mathcal{L}$  = a selection of d.o.f.'s
- ▶ 'small' **phase** space  $\mathcal{M}_\eta$


Coarse graining:

- ▶ projections  $\pi_{\eta' \rightarrow \eta}$  for  $\eta \preccurlyeq \eta'$
- ▶ lifting observables  
 $A_\eta \circ \pi_{\eta' \rightarrow \eta} =: A_{\eta'}$   
↳ compatible with the Poisson brackets
- ▶ **preferred** factorization  
 $\mathcal{M}_{\eta'} \approx \tilde{\mathcal{M}}_{\eta' \rightarrow \eta} \times \mathcal{M}_\eta$

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective Formalism for QFT

Coarse-graining between Quantum State Spaces



Quantization:

- ▶ classical factorizations  
~~~  $\otimes$ -factorizations  
 $\mathcal{H}_{\eta'} \approx \mathcal{H}_{\eta' \rightarrow \eta} \otimes \mathcal{H}_\eta$
- ▶ **unambiguous**  $\tilde{\mathcal{H}}_{\eta' \rightarrow \eta} \approx \mathcal{H}_{\eta' \rightarrow \eta}$

Projective families  $(\rho_\eta)_{\eta \in \mathcal{L}}$ :

- ▶  $\rho_\eta$  density matrix on  $\mathcal{H}_\eta$
- ▶  $\rho_\eta = \text{Tr}_{\mathcal{H}_{\eta' \rightarrow \eta}} \rho_{\eta'} = \text{Tr}_{\tilde{\mathcal{H}}_{\eta' \rightarrow \eta}} \rho_{\eta'}$


Ex: linear QFT

~~~ **polarization-independant**  
quantum state space

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective Formalism for QFT

Coarse-graining between Quantum State Spaces



Quantization:

- ▶ classical factorizations  
~~~  $\otimes$ -factorizations  
 $\mathcal{H}_{\eta'} \approx \mathcal{H}_{\eta' \rightarrow \eta} \otimes \mathcal{H}_{\eta}$
- ▶ **unambiguous**  $\tilde{\mathcal{H}}_{\eta' \rightarrow \eta} \approx \mathcal{H}_{\eta' \rightarrow \eta}$

Projective families  $(\rho_{\eta})_{\eta \in \mathcal{L}}$ :

- ▶  $\rho_{\eta}$  density matrix on  $\mathcal{H}_{\eta}$
- ▶  $\rho_{\eta} = \text{Tr}_{\mathcal{H}_{\eta' \rightarrow \eta}} \rho_{\eta'} = \text{Tr}_{\tilde{\mathcal{H}}_{\eta' \rightarrow \eta}} \rho_{\eta'}$


Ex: linear QFT

~~~ **polarization-independant**  
quantum state space

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective Formalism for QFT

Coarse-graining between Quantum State Spaces



Quantization:

- ▶ classical factorizations  
~~>  $\otimes$ -factorizations  
 $\mathcal{H}_{\eta'} \approx \mathcal{H}_{\eta' \rightarrow \eta} \otimes \mathcal{H}_\eta$
- ▶ **unambiguous**  $\tilde{\mathcal{H}}_{\eta' \rightarrow \eta} \approx \mathcal{H}_{\eta' \rightarrow \eta}$

Projective families  $(\rho_\eta)_{\eta \in \mathcal{L}}$ :

- ▶  $\rho_\eta$  density matrix on  $\mathcal{H}_\eta$
- ▶  $\rho_\eta = \text{Tr}_{\mathcal{H}_{\eta' \rightarrow \eta}} \rho_{\eta'} = \text{Tr}_{\tilde{\mathcal{H}}_{\eta' \rightarrow \eta}} \rho_{\eta'}$

Ex: linear QFT

~~> **polarization-independant**  
quantum state space

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective Formalism for QFT

Natural Embedding of Arbitrary Vacuum Sectors

$$\boxed{\begin{aligned}\mathcal{H}_{\eta'} &\approx \mathcal{H}_{\eta' \rightarrow \eta} \otimes \mathcal{H}_\eta \\ \rho_{\eta'} &\end{aligned}}$$

↓  
partial  
trace

$$\boxed{\begin{aligned}\mathcal{H}_\eta & \\ \rho_\eta = \text{Tr}_{\eta' \rightarrow \eta} \rho_{\eta'} &\end{aligned}}$$

$$\eta \preccurlyeq \eta' \in \mathcal{L}$$

Inductive limit  $\mathcal{F}_\emptyset$ :

- ▶ choice of **vacuum**

$$\forall \eta \preccurlyeq \eta', |\emptyset_{\eta' \rightarrow \eta}\rangle \in \mathcal{H}_{\eta' \rightarrow \eta}$$

$$\rightsquigarrow \text{injection } \iota_{\eta' \leftarrow \eta} : \mathcal{H}_\eta \rightarrow \mathcal{H}_{\eta'}$$

- ▶ ex: Fock representation,  
LQG state space,...

Mapping a density matrix  $\sigma$  on  $\mathcal{F}_\emptyset$   
to a projective state  $(\rho_\eta)_\eta$ :

- ▶ this mapping is **injective**
- ▶ its image can be characterized

[GNS Representation: Gelfand, Naimark, Segal]

Projective State Spaces for QG (S. Lanéry)

└ Projective State Spaces


└ Embedding of Fock Repr.

arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)

QProjStSp InjFock InjAL FockProjQ

# Projective Formalism for QFT

Natural Embedding of Arbitrary Vacuum Sectors



Inductive limit  $\mathcal{F}_\emptyset$ :

- choice of **vacuum**

$$\forall \eta \preccurlyeq \eta', |\emptyset_{\eta' \rightarrow \eta}\rangle \in \mathcal{H}_{\eta' \rightarrow \eta}$$

$$\rightsquigarrow \text{injection } \iota_{\eta' \leftarrow \eta} : \mathcal{H}_\eta \rightarrow \mathcal{H}_{\eta'}$$

- ex: Fock representation,  
LQG state space,...

Mapping a density matrix  $\sigma$  on  $\mathcal{F}_\emptyset$  to a projective state  $(\rho_\eta)_\eta$ :

- this mapping is **injective**
- its image can be characterized

[GNS Representation: Gelfand, Naimark, Segal]

Projective State Spaces for QG (S. Lanéry)

└ Projective State Spaces

└ Embedding of Fock Repr.

arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)

QProjStSp InjFock InjAL FockProjQ

# Contents

## Projective Formalism for QFT

### Universal Selections of Degrees of Freedom

    Restriction to a Cofinal Sequence of Labels

    (Semi-classical) States on a Sequence of Labels


    Restriction to a Quasi-Cofinal Sequence of Labels

    Universality & Symmetries

## Application to Loop Quantum Gravity

# Universal Selections of Degrees of Freedom

Restriction to a Cofinal Sequence of Labels



Sequence  $(\kappa_n)_{n \geq 1}$ :


- increasing:  $\kappa_n \preccurlyeq \kappa_{n+1}$
- **cofinal**:  $\forall \eta \in \mathcal{L}, \exists n / \eta \preccurlyeq \kappa_n$

Bijective mapping  $(\rho_{\kappa_n})_n \leftrightarrow (\rho_\eta)_{\eta \in \mathcal{L}}$ :

- $\rho_\eta = \text{Tr}_{\kappa_n \rightarrow \eta} \rho_{\kappa_n}$  for some  $\kappa_n \succcurlyeq \eta$
- projective quantum state spaces are **robust** (by contrast eg. to infinite tensor products)

# Universal Selections of Degrees of Freedom

Restriction to a Cofinal Sequence of Labels



Sequence  $(\kappa_n)_{n \geq 1}$ :


- increasing:  $\kappa_n \preccurlyeq \kappa_{n+1}$
- **cofinal**:  $\forall \eta \in \mathcal{L}, \exists n / \eta \preccurlyeq \kappa_n$

Bijective mapping  $(\rho_{\kappa_n})_n \leftrightarrow (\rho_\eta)_{\eta \in \mathcal{L}}$ :

- $\rho_\eta = \text{Tr}_{\kappa_n \rightarrow \eta} \rho_{\kappa_n}$  for some  $\kappa_n \succcurlyeq \eta$
- projective quantum state spaces are **robust** (by contrast eg. to infinite tensor products)

# Universal Selections of Degrees of Freedom

Restriction to a Cofinal Sequence of Labels



Sequence  $(\kappa_n)_{n \geq 1}$ :

- increasing:  $\kappa_n \preccurlyeq \kappa_{n+1}$
- **cofinal**:  $\forall \eta \in \mathcal{L}, \exists n / \eta \preccurlyeq \kappa_n$

Bijective mapping  $(\rho_{\kappa_n})_n \leftrightarrow (\rho_\eta)_{\eta \in \mathcal{L}}$ :

- $\rho_\eta = \text{Tr}_{\kappa_n \rightarrow \eta} \rho_{\kappa_n}$  for some  $\kappa_n \succcurlyeq \eta$
- projective quantum state spaces are **robust** (by contrast eg. to infinite tensor products)

# Universal Selections of Degrees of Freedom

(Semi-classical) States on a Sequence of Labels

Semi-classical states on an increasing sequence  $(\kappa_n)_{n \geq 1}$ :

- ▶ first at **macroscopic** scale
- ▶ then, step by step, on complementary d.o.f.'s

$$\kappa_{n+1} \rightarrow \kappa_n$$

~~ saturating uncertainty relations, given already chosen macro scale behavior

$$\mathcal{H}_{\kappa_1}$$

$$\rho_{\kappa_1} = |\zeta_{\kappa_1} \times \zeta_{\kappa_1}|$$

$$\kappa_1 \in \mathcal{L}$$

More generally: systematic, **recursive** construction of all projective states  $(\rho_{\kappa_n})_{n \geq 1}$

[See also: Giesel & Thiemann '06; Oriti, Pereira & Sindoni '12]

# Universal Selections of Degrees of Freedom

(Semi-classical) States on a Sequence of Labels

$$\mathcal{H}_{\kappa_{n+1}} \approx \mathcal{H}_{\kappa_{n+1} \rightarrow \kappa_n} \otimes \mathcal{H}_{\kappa_n}$$

$$\rho_{\kappa_{n+1}} = |\zeta_{\kappa_{n+1} \rightarrow \kappa_n} \times \zeta_{\kappa_{n+1} \rightarrow \kappa_n}| \otimes \rho_{\kappa_n}$$

$$\mathcal{H}_{\kappa_n}$$

$$\rho_{\kappa_n} = |\zeta_{\kappa_n} \times \zeta_{\kappa_n}|$$

$$\kappa_n \preccurlyeq \kappa_{n+1} \in \mathcal{L}$$

Semi-classical states on an increasing sequence  $(\kappa_n)_{n \geq 1}$ :

- ▶ first at **macroscopic** scale
- ▶ then, step by step, on complementary d.o.f.'s

$$\kappa_{n+1} \rightarrow \kappa_n$$

$\rightsquigarrow$  saturating uncertainty relations, given already chosen macro scale behavior

More generally: systematic, **recursive** construction of all projective states  $(\rho_{\kappa_n})_{n \geq 1}$

[See also: Giesel & Thiemann '06; Oriti, Pereira & Sindoni '12]

# Universal Selections of Degrees of Freedom

(Semi-classical) States on a Sequence of Labels

$$\mathcal{H}_{\kappa_{n+1}} \approx \mathcal{H}_{\kappa_{n+1} \rightarrow \kappa_n} \otimes \mathcal{H}_{\kappa_n}$$
$$\rho_{\kappa_{n+1}} = |\zeta_{\kappa_{n+1} \rightarrow \kappa_n} \times \zeta_{\kappa_{n+1} \rightarrow \kappa_n}| \otimes \rho_{\kappa_n}$$

partial trace  $\downarrow$

$$\text{Tr}_{\kappa_{n+1} \rightarrow \kappa_n}$$

$$\mathcal{H}_{\kappa_n}$$
$$\rho_{\kappa_n} = |\zeta_{\kappa_n} \times \zeta_{\kappa_n}|$$

$$\kappa_n \preccurlyeq \kappa_{n+1} \in \mathcal{L}$$

Semi-classical states on an increasing sequence  $(\kappa_n)_{n \geq 1}$ :


- ▶ first at **macroscopic** scale
- ▶ then, step by step, on complementary d.o.f.'s  
 $\kappa_{n+1} \rightarrow \kappa_n$   
~~~ saturating uncertainty relations, given already chosen macro scale behavior

More generally: systematic, **recursive** construction of all projective states  $(\rho_{\kappa_n})_{n \geq 1}$

[See also: Giesel & Thiemann '06; Oriti, Pereira & Sindoni '12]

# Universal Selections of Degrees of Freedom

Restriction to a Quasi-Cofinal Sequence of Labels



Continuum of d.o.f.'s:

- ▶ abstraction for math convenience
- ▶ **dense, countable** sub-algebra of observables is enough  $\rightsquigarrow$  **but:** ensure **universality** and respect **symmetries** of the theory


**Quasi-cofinal** increasing sequence:

$$\forall \eta \in \mathcal{L}, \exists n \geq 1 /$$

- ▶  $\tau \cdot \eta \preccurlyeq \kappa_n$  (with  $\tau$  small deformation)
- ▶  $\forall \kappa \preccurlyeq \eta, \kappa_m, \tau \cdot \kappa \stackrel{!}{=} \kappa$   
(for any  $m \geq 1$ )

# Universal Selections of Degrees of Freedom

Restriction to a Quasi-Cofinal Sequence of Labels



Continuum of d.o.f.'s:

- abstraction for math convenience
- **dense, countable** sub-algebra of observables is enough  $\rightsquigarrow$  **but:** ensure **universality** and respect **symmetries** of the theory


**Quasi-cofinal** increasing sequence:

$$\forall \eta \in \mathcal{L}, \exists n \geq 1 /$$

- $\tau \cdot \eta \preccurlyeq \kappa_n$  (with  $\tau$  small deformation)
- $\forall \kappa \preccurlyeq \eta, \kappa_m, \tau \cdot \kappa \stackrel{!}{=} \kappa$   
(for any  $m \geq 1$ )

# Universal Selections of Degrees of Freedom

Restriction to a Quasi-Cofinal Sequence of Labels



Continuum of d.o.f.'s:

- ▶ abstraction for math convenience
- ▶ **dense, countable** sub-algebra of observables is enough  $\rightsquigarrow$  **but:** ensure **universality** and respect **symmetries** of the theory


**Quasi-cofinal** increasing sequence:

$$\forall \eta \in \mathcal{L}, \exists n \geq 1 /$$

- ▶  $\tau \cdot \eta \preccurlyeq \kappa_n$  (with  $\tau$  small deformation)
- ▶  $\forall \kappa \preccurlyeq \eta, \kappa_m, \tau \cdot \kappa \stackrel{!}{=} \kappa$   
(for any  $m \geq 1$ )

# Universal Selections of Degrees of Freedom

Restriction to a Quasi-Cofinal Sequence of Labels



Continuum of d.o.f.'s:

- abstraction for math convenience
- **dense, countable** sub-algebra of observables is enough  $\rightsquigarrow$  **but:** ensure **universality** and respect **symmetries** of the theory


**Quasi-cofinal** increasing sequence:

$$\forall \eta \in \mathcal{L}, \exists n \geq 1 /$$

- $\tau \cdot \eta \preccurlyeq \kappa_n$  (with  $\tau$  small deformation)
- $\forall \kappa \preccurlyeq \eta, \kappa_m, \tau \cdot \kappa \stackrel{!}{=} \kappa$   
(for any  $m \geq 1$ )

# Universal Selections of Degrees of Freedom

Restriction to a Quasi-Cofinal Sequence of Labels



Continuum of d.o.f.'s:

- abstraction for math convenience
- **dense, countable** sub-algebra of observables is enough  $\rightsquigarrow$  **but:** ensure **universality** and respect **symmetries** of the theory


**Quasi-cofinal** increasing sequence:

$$\forall \eta \in \mathcal{L}, \exists n \geq 1 /$$

- $\tau \cdot \eta \preccurlyeq \kappa_n$  (with  $\tau$  small deformation)
- $\forall \kappa \preccurlyeq \eta, \kappa_m, \tau \cdot \kappa \stackrel{!}{=} \kappa$   
(for any  $m \geq 1$ )

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta_{m_k} &\preccurlyeq \tau \cdot \kappa_{n_k} \\ \blacktriangleright \tau \cdot \kappa_{n_k} &\preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow (s \cdot \eta_m)_m &\sim (s \cdot \eta_{m_k})_k \sim \\ (\tau \cdot \kappa_{n_k})_k &\sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of  
the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta_{m_k} &\preccurlyeq \tau \cdot \kappa_{n_k} \\ \blacktriangleright \tau \cdot \kappa_{n_k} &\preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow (s \cdot \eta_m)_m &\sim (s \cdot \eta_{m_k})_k \sim \\ (\tau \cdot \kappa_{n_k})_k &\sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of  
the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta m_k &\preccurlyeq \tau \cdot \kappa n_k \\ \blacktriangleright \tau \cdot \kappa n_k &\preccurlyeq s \cdot \eta m_{k+1} \\ \Rightarrow (s \cdot \eta m)_m &\sim (s \cdot \eta m_k)_k \sim \\ (\tau \cdot \kappa n)_k &\sim (\tau \cdot \kappa n)_n \end{aligned}$$

Approximating symmetries of  
the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} &\blacktriangleright s \cdot \eta_{m_k} \preccurlyeq \tau \cdot \kappa_{n_k} \\ &\blacktriangleright \tau \cdot \kappa_{n_k} \preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow & (s \cdot \eta_m)_m \sim (s \cdot \eta_{m_k})_k \sim \\ & (\tau \cdot \kappa_{n_k})_k \sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of the theory (eg. diffeos):

$$\begin{aligned} &\blacktriangleright (\Phi \cdot \kappa_n)_n =: (\eta_m)_m \\ &\blacktriangleright (s \cdot \eta_m)_m \sim (\tau \cdot \kappa_n)_n \\ \Rightarrow & (\tilde{\Phi} \cdot \kappa_n)_n \sim (\kappa_n)_n \\ & \text{(with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta_{m_k} &\preccurlyeq \tau \cdot \kappa_{n_k} \\ \blacktriangleright \tau \cdot \kappa_{n_k} &\preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow (s \cdot \eta_m)_m &\sim (s \cdot \eta_{m_k})_k \sim \\ (\tau \cdot \kappa_{n_k})_k &\sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta_{m_k} &\preccurlyeq \tau \cdot \kappa_{n_k} \\ \blacktriangleright \tau \cdot \kappa_{n_k} &\preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow (s \cdot \eta_m)_m &\sim (s \cdot \eta_{m_k})_k \sim \\ (\tau \cdot \kappa_{n_k})_k &\sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta_{m_k} &\preccurlyeq \tau \cdot \kappa_{n_k} \\ \blacktriangleright \tau \cdot \kappa_{n_k} &\preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow (s \cdot \eta_m)_m &\sim (s \cdot \eta_{m_k})_k \sim \\ (\tau \cdot \kappa_{n_k})_k &\sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of  
the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} &\blacktriangleright s \cdot \eta_{m_k} \preccurlyeq \tau \cdot \kappa_{n_k} \\ &\blacktriangleright \tau \cdot \kappa_{n_k} \preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow & (s \cdot \eta_m)_m \sim (s \cdot \eta_{m_k})_k \sim \\ & (\tau \cdot \kappa_{n_k})_k \sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of the theory (eg. diffeos):

$$\begin{aligned} &\blacktriangleright (\Phi \cdot \kappa_n)_n =: (\eta_m)_m \\ &\blacktriangleright (s \cdot \eta_m)_m \sim (\tau \cdot \kappa_n)_n \\ \Rightarrow & (\tilde{\Phi} \cdot \kappa_n)_n \sim (\kappa_n)_n \\ & (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta_{m_k} &\preccurlyeq \tau \cdot \kappa_{n_k} \\ \blacktriangleright \tau \cdot \kappa_{n_k} &\preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow (s \cdot \eta_m)_m &\sim (s \cdot \eta_{m_k})_k \sim \\ (\tau \cdot \kappa_{n_k})_k &\sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta_{m_k} &\preccurlyeq \tau \cdot \kappa_{n_k} \\ \blacktriangleright \tau \cdot \kappa_{n_k} &\preccurlyeq s \cdot \eta_{m_{k+1}} \\ \Rightarrow (s \cdot \eta_m)_m &\sim (s \cdot \eta_{m_k})_k \sim \\ (\tau \cdot \kappa_{n_k})_k &\sim (\tau \cdot \kappa_n)_n \end{aligned}$$

Approximating symmetries of the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

## Universality & Symmetries



Subsequences + deformations /


$$\begin{aligned} \blacktriangleright s \cdot \eta m_k &\preccurlyeq \tau \cdot \kappa n_k \\ \blacktriangleright \tau \cdot \kappa n_k &\preccurlyeq s \cdot \eta m_{k+1} \\ \Rightarrow (s \cdot \eta m)_m &\sim (s \cdot \eta m_k)_k \sim \\ (\tau \cdot \kappa n_k)_k &\sim (\tau \cdot \kappa n)_n \end{aligned}$$

Approximating symmetries of the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Universal Selections of Degrees of Freedom

Universality & Symmetries



Subsequences + deformations /

$$\begin{aligned} \blacktriangleright s \cdot \eta m_k &\preccurlyeq \tau \cdot \kappa n_k \\ \blacktriangleright \tau \cdot \kappa n_k &\preccurlyeq s \cdot \eta m_{k+1} \\ \Rightarrow (s \cdot \eta m)_m &\sim (s \cdot \eta m_k)_k \sim \\ (\tau \cdot \kappa n_k)_k &\sim (\tau \cdot \kappa n)_n \end{aligned}$$

Approximating symmetries of  
the theory (eg. diffeos):

$$\begin{aligned} \blacktriangleright (\Phi \cdot \kappa_n)_n &=: (\eta_m)_m \\ \blacktriangleright (s \cdot \eta_m)_m &\sim (\tau \cdot \kappa_n)_n \\ \Rightarrow (\tilde{\Phi} \cdot \kappa_n)_n &\sim (\kappa_n)_n \\ (\text{with } \tilde{\Phi} := \tau^{-1} \circ s \circ \Phi) \end{aligned}$$

# Contents

Projective Formalism for QFT

Universal Selections of Degrees of Freedom

Application to Loop Quantum Gravity


Projective State Space for LQG

Relation to the AL Hilbert Space

Fractal Sequences of Labels

# Application to Loop Quantum Gravity

## Projective State Space for LQG



The labels:

- ▶ a graph = holonomies
- ▶ a set of surfaces dual to this graph = conjugate fluxes
- ▶  $\mathcal{H}_\eta := L_2(G^n, d\mu_{\text{Haar}})$



The coarse-grainings:

- ▶  $G^n \approx G^m \times G^{n-m}$
- ▶ kept holonomies  
~~ prescribes the factor  $G^m$
- ▶ kept fluxes ~~  
complementary factor  $G^{n-m}$

[Holonomy-flux algebra: Ashtekar, Gambini, Isham, Lewandowski, Pullin, Rovelli, Smolin,...]

# Application to Loop Quantum Gravity

## Projective State Space for LQG



The labels:

- ▶ a graph = holonomies
- ▶ a set of surfaces dual to this graph = conjugate fluxes
- ▶  $\mathcal{H}_\eta := L_2(G^n, d\mu_{\text{Haar}})$



The coarse-grainings:

- ▶  $G^n \approx G^m \times G^{n-m}$
- ▶ kept holonomies  
 $\rightsquigarrow$  prescribes the factor  $G^m$
- ▶ kept fluxes  $\rightsquigarrow$   
complementary factor  $G^{n-m}$

[Holonomy-flux algebra: Ashtekar, Gambini, Isham, Lewandowski, Pullin, Rovelli, Smolin,...]

# Application to Loop Quantum Gravity

## Projective State Space for LQG



The labels:

- ▶ a graph = holonomies
- ▶ a set of surfaces dual to this graph = conjugate fluxes
- ▶  $\mathcal{H}_\eta := L_2(G^n, d\mu_{\text{Haar}})$



The coarse-grainings:

- ▶  $G^n \approx G^m \times G^{n-m}$
- ▶ kept holonomies  
 $\rightsquigarrow$  prescribes the factor  $G^m$
- ▶ kept fluxes  $\rightsquigarrow$   
complementary factor  $G^{n-m}$

[Holonomy-flux algebra: Ashtekar, Gambini, Isham, Lewandowski, Pullin, Rovelli, Smolin,...]

# Application to Loop Quantum Gravity

## Projective State Space for LQG



The labels:

- ▶ a graph = holonomies
- ▶ a set of surfaces dual to this graph = conjugate fluxes
- ▶  $\mathcal{H}_\eta := L_2(G^n, d\mu_{\text{Haar}})$



The coarse-grainings:

- ▶  $G^n \approx G^m \times G^{n-m}$
- ▶ kept holonomies  
 $\rightsquigarrow$  prescribes the factor  $G^m$
- ▶ kept fluxes  $\rightsquigarrow$   
complementary factor  $G^{n-m}$

[Holonomy-flux algebra: Ashtekar, Gambini, Isham, Lewandowski, Pullin, Rovelli, Smolin,...]

# Application to Loop Quantum Gravity

## Projective State Space for LQG



The labels:

- ▶ a graph = holonomies
- ▶ a set of surfaces dual to this graph = conjugate fluxes
- ▶  $\mathcal{H}_\eta := L_2(G^n, d\mu_{\text{Haar}})$



The coarse-grainings:

- ▶  $G^n \approx G^m \times G^{n-m}$
- ▶ kept holonomies  
~~ prescribes the factor  $G^m$
- ▶ kept fluxes ~~  
complementary factor  $G^{n-m}$

[Holonomy-flux algebra: Ashtekar, Gambini, Isham, Lewandowski, Pullin, Rovelli, Smolin,...]

# Application to Loop Quantum Gravity

## Projective State Space for LQG



The labels:

- ▶ a graph = holonomies
- ▶ a set of surfaces dual to this graph = conjugate fluxes
- ▶  $\mathcal{H}_\eta := L_2(G^n, d\mu_{\text{Haar}})$



The coarse-grainings:

- ▶  $G^n \approx G^m \times G^{n-m}$
- ▶ kept holonomies  
 $\rightsquigarrow$  prescribes the factor  $G^m$
- ▶ kept fluxes  $\rightsquigarrow$   
complementary factor  $G^{n-m}$

[Holonomy-flux algebra: Ashtekar, Gambini, Isham, Lewandowski, Pullin, Rovelli, Smolin,...]

# Application to Loop Quantum Gravity

## Projective State Space for LQG



The labels:

- ▶ a graph = holonomies
- ▶ a set of surfaces dual to this graph = conjugate fluxes
- ▶  $\mathcal{H}_\eta := L_2(G^n, d\mu_{\text{Haar}})$


The coarse-grainings:

- ▶  $G^n \approx G^m \times G^{n-m}$
- ▶ kept holonomies  
~~ prescribes the factor  $G^m$
- ▶ kept fluxes ~~  
complementary factor  $G^{n-m}$


[Holonomy-flux algebra: Ashtekar, Gambini, Isham, Lewandowski, Pullin, Rovelli, Smolin,...]

# Application to Loop Quantum Gravity

## Relation to the AL Hilbert Space



$$\psi_\eta(h_1, h_2) = \psi_{\eta''}(h_1, h_2, \dots)$$



Inductive limit  $\mathcal{H}_{AL}$ :

- ▶ AL vacuum:  $\emptyset_{\eta' \rightarrow \eta} \equiv 1 \in \mathcal{H}_{\eta' \rightarrow \eta} = L_2(G^{n-m})$
- ▶ injection  $\mathcal{H}_\eta \rightarrow \mathcal{H}_{\eta'}$  only depends on the **graphs**

Projective state space extends AL state space:

- ▶ holonomies and fluxes on the same footing
- ▶ first step toward better semi-classical states

[LQG Hilbert space: Ashtekar, Isham, Lewandowski,...]

Projective State Spaces for QG (S. Lanéry)

└ Projective LQG


└ AL Hilbert Space

arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)


LQG ALHilb InjGI FactVsProj ALFacto

# Application to Loop Quantum Gravity

## Relation to the AL Hilbert Space



$$\psi_\eta(h_1, h_2) = \psi_{\eta''}(h_1, h_2, \dots) = \psi_{\eta'}(h_1, h_2)$$



Inductive limit  $\mathcal{H}_{\text{AL}}$ :

- ▶ AL vacuum:  $\emptyset_{\eta' \rightarrow \eta} \equiv 1 \in \mathcal{H}_{\eta' \rightarrow \eta} = L_2(G^{n-m})$
- ▶ injection  $\mathcal{H}_\eta \rightarrow \mathcal{H}_{\eta'}$  only depends on the **graphs**

Projective state space extends AL state space:

- ▶ holonomies and fluxes on the same footing
- ▶ first step toward better semi-classical states

[LQG Hilbert space: Ashtekar, Isham, Lewandowski,...]

Projective State Spaces for QG (S. Lanéry)

└ Projective LQG


└ AL Hilbert Space

arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)

LQG ALHilb InjGI FactVsProj ALFacto

# Application to Loop Quantum Gravity

## Fractal Sequences of Labels



**Fractal**-like sequence of labels:

- ▶ **universality** (equivalent state spaces from  $\neq$  sequences)
- ▶ **background independence** (approx any diffeomorphism by fractal-stabilizing morphism)


Viz. semi-analytic structure:

- ▶  $\neq$  semi-analytic structures related by small deformation
- ▶ any diffeo can be approx by semi-analytic one

[See also: Renteln & Smolin, Gambini & Pullin, Corichi & Zapata, Loll,...]

# Application to Loop Quantum Gravity

## Fractal Sequences of Labels



**Fractal**-like sequence of labels:

- ▶ **universality** (equivalent state spaces from  $\neq$  sequences)
- ▶ **background independence** (approx any diffeomorphism by fractal-stabilizing morphism)


Viz. semi-analytic structure:

- ▶  $\neq$  semi-analytic structures related by small deformation
- ▶ any diffeo can be approx by semi-analytic one

[See also: Renteln & Smolin, Gambini & Pullin, Corichi & Zapata, Loll,...]

# Application to Loop Quantum Gravity

## Fractal Sequences of Labels



$$\kappa_1 \preccurlyeq \kappa_2 \preccurlyeq \kappa_3$$

**Fractal-like sequence of labels:**

- ▶ **universality** (equivalent state spaces from  $\neq$  sequences)
- ▶ **background independence** (approx any diffeomorphism by fractal-stabilizing morphism)

Viz. semi-analytic structure:

- ▶  $\neq$  semi-analytic structures related by small deformation
- ▶ any diffeo can be approx by semi-analytic one

[See also: Renteln & Smolin, Gambini & Pullin, Corichi & Zapata, Loll,...]

Projective State Spaces for QG (S. Lanéry)

└ Projective LQG

└ Fractals

arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)

QuasiCfnl QuasiCfnlUniv Fractal1D FractalUniv

# Summary: Comparison with other State Spaces

|                                        | analytic<br>holo. & fluxes<br>$\mathcal{H}_{\text{AL}}$ | infinite<br>graph<br>AQG | fractal-based<br>sub-algebra<br>induc. limit | proj. st.<br>space | ITP |
|----------------------------------------|---------------------------------------------------------|--------------------------|----------------------------------------------|--------------------|-----|
| Hilbert space                          | ✓                                                       | ✗                        | ✓                                            | ✓                  | ✗   |
| Separability                           | ✗                                                       | "✗"                      | ✗                                            | ✓                  | "✓" |
| Universality +<br>Action of diffeos.   | ✓                                                       | ✓                        | ✗                                            | ✓                  | ✓   |
| Solving Gauss &<br>diffeo. constraints | ✓                                                       | ✗                        | ✓                                            | ✓                  | ?   |
| Non-compact $G$                        | ✗                                                       | ?                        | ✓                                            | ✗                  | ✓   |
| Untruncated<br>semi-cl. states         | ✗                                                       | ✗                        | ✓                                            | ✗                  | ✓   |

[ITP: von Neumann '39; AQG: Giesel & Thiemann '06,...]


# Outlook

Applications of the thus constructed state space:

- ▶ study of the **semi-classical regime** of loop quantum gravity
- ▶ derivation of **symmetry reduced models** (LQC, spherically symmetric LQG,...) from the full theory [Engle '07]

Implementation of the **constraints**:

- ▶ general prescriptions to deal with constraints in the context of projective state spaces: in particular, relations with **covariant approaches**, renormalization...
- ▶ **Gauss & diffeomorphism** constraints: somewhat easier when working on fractal sequence of labels
- ▶ dynamics...



Thank you!

# Contents

## Extra Slides

Projective State Spaces

Relation to Other Constructions

Implementation of Gauge Constraints and Dynamics


Basics of (Loop) Quantum Gravity

Projective State Space for the Holonomy-Flux Algebra

Obstruction to the Construction of Semi-classical States

# Classical Projective State Spaces

## Projective Systems of Phase Spaces



$$\eta \preccurlyeq \eta' \preccurlyeq \eta'' \in \mathcal{L}$$

Collection of partial theories:

- ▶ label set  $\mathcal{L}$ ,  $\preccurlyeq$
- ▶  $\eta \in \mathcal{L}$  = a selection of d.o.f.'s
- ▶ 'small' phase spaces  $\mathcal{M}_\eta$

Ensuring **consistency**:

- ▶ projections  $\pi_{\eta' \rightarrow \eta}$  for  $\eta \preccurlyeq \eta'$
- ▶ mounting observables:  
 $O_{\eta'} = O_\eta \circ \pi_{\eta' \rightarrow \eta}$
- ▶ 3-spaces-consistency  
→ projective system

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective State Spaces

## Projections and Factorizations

$$\pi : \mathcal{M} \rightarrow \widetilde{\mathcal{M}}$$

$q_1, \dots, q_n$   
 $p_1, \dots, p_n;$



$\tilde{q}_1, \dots, \tilde{q}_m$   
 $\tilde{p}_1, \dots, \tilde{p}_m;$

# Projective State Spaces

## Projections and Factorizations

$$\pi : \mathcal{M} \rightarrow \widetilde{\mathcal{M}}$$

$$\mathcal{M} \approx \widetilde{\mathcal{M}} \times \underline{\mathcal{M}}$$

$q_1, \dots, q_n$   
 $p_1, \dots, p_n;$



$\tilde{q}_1, \dots, \tilde{q}_m$   
 $\tilde{p}_1, \dots, \tilde{p}_m;$

$q_1, \dots, q_n$   
 $p_1, \dots, p_n;$



$\tilde{q}_1, \dots, \tilde{q}_m, \underline{q}_{m+1}, \dots, \underline{q}_n$   
 $\tilde{p}_1, \dots, \tilde{p}_m, \underline{p}_{m+1}, \dots, \underline{p}_n;$

# Projective State Spaces

## Projections and Factorizations

$$\pi : \mathcal{M} \rightarrow \widetilde{\mathcal{M}}$$

$$q_1, \dots, q_n \\ p_1, \dots, p_n;$$



$$\widetilde{q}_1, \dots, \widetilde{q}_m \\ \widetilde{p}_1, \dots, \widetilde{p}_m;$$

$$\mathcal{M} \approx \widetilde{\mathcal{M}} \times \underline{\mathcal{M}}$$

$$q_1, \dots, q_n \\ p_1, \dots, p_n;$$



$$\widetilde{q}_1, \dots, \widetilde{q}_m, \underline{q}_{m+1}, \dots, \underline{q}_n \\ \widetilde{p}_1, \dots, \widetilde{p}_m, \underline{p}_{m+1}, \dots, \underline{p}_n;$$

$$\mathcal{C} \approx \widetilde{\mathcal{C}} \times \underline{\mathcal{C}}$$

$$q_1, \dots, q_n$$



$$\widetilde{q}_1, \dots, \widetilde{q}_m, \underline{q}_{m+1}, \dots, \underline{q}_n$$

# Projective State Spaces

## Projections and Factorizations

$$\pi : \mathcal{M} \rightarrow \widetilde{\mathcal{M}}$$

$$\mathcal{M} \approx \widetilde{\mathcal{M}} \times \underline{\mathcal{M}}$$

$$\mathcal{C} \approx \widetilde{\mathcal{C}} \times \underline{\mathcal{C}}$$

$q_1, \dots, q_n$   
 $p_1, \dots, p_n;$



$\tilde{q}_1, \dots, \tilde{q}_m$   
 $\tilde{p}_1, \dots, \tilde{p}_m;$

$q_1, \dots, q_n$   
 $p_1, \dots, p_n;$



$\tilde{q}_1, \dots, \tilde{q}_m, \underline{q}_{m+1}, \dots, \underline{q}_n$   
 $\tilde{p}_1, \dots, \tilde{p}_m, \underline{p}_{m+1}, \dots, \underline{p}_n;$

$q_1, \dots, q_n$



$\tilde{q}_1, \dots, \tilde{q}_m, \underline{q}_{m+1}, \dots, \underline{q}_n$

---

$\neq$

---

$$\tau : \mathcal{C} \rightarrow \widetilde{\mathcal{C}}$$

$q_1, \dots, q_n$




$\tilde{q}_1, \dots, \tilde{q}_m$

# Classical Projective State Spaces

Example: Second Quantization

$$I' \approx (I^\perp \cap I') \times I$$




First quantized theory:

- ▶ Hilbert space  $\mathcal{H}_{\text{one particule}}$   
 $\omega_{\text{symplect}} := 2 \operatorname{Im} \langle \cdot, \cdot \rangle$
- ▶  $a_e : \psi \mapsto \langle e, \psi \rangle$
- ▶  $\{a_e, a_f^*\} = i \langle e, f \rangle$

Projective system:

- ▶ labels:  $I \subset \mathcal{H}_{\text{one particule}}$  with  $\dim I < \infty$
- ▶ ‘small’ phase space:  $\mathcal{M}_I := I$
- ▶  $\pi_{I' \rightarrow I}$  orthog. projection on  $I \subset I'$
- ▶  $\exists!$  factorization:  $I' \approx (I^\perp \cap I') \times I$

# Quantum Projective State Spaces



Modeled on special case:


- ▶ classical factorizations  
 $\mathcal{M}_{\eta'} \approx \mathcal{M}_{\eta' \rightarrow \eta} \times \mathcal{M}_\eta$
- ▶ 3-spaces-consistency  
 $\mathcal{M}_{\eta'' \rightarrow \eta} \approx \mathcal{M}_{\eta'' \rightarrow \eta'} \times \mathcal{M}_{\eta' \rightarrow \eta}$
- ▶ quantization  
 $\rightsquigarrow \otimes\text{-factorizations}$

Projective families  $(\rho_\eta)_{\eta \in \mathcal{L}}$ :

- ▶  $\rho_\eta$  density matrix on  $\mathcal{H}_\eta$
- ▶  $\mathrm{Tr}_{\mathcal{H}_{\eta' \rightarrow \eta}} \rho_{\eta'} = \rho_\eta$

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Quantum Projective State Spaces



$$\eta \preccurlyeq \eta' \preccurlyeq \eta'' \in \mathcal{L}$$

Modeled on special case:

- ▶ classical factorizations  
 $\mathcal{M}_{\eta'} \approx \mathcal{M}_{\eta' \rightarrow \eta} \times \mathcal{M}_\eta$
- ▶ 3-spaces-consistency  
 $\mathcal{M}_{\eta'' \rightarrow \eta} \approx \mathcal{M}_{\eta'' \rightarrow \eta'} \times \mathcal{M}_{\eta' \rightarrow \eta}$
- ▶ quantization  
 $\rightsquigarrow \otimes\text{-factorizations}$

Projective families  $(\rho_\eta)_{\eta \in \mathcal{L}}$ :

- ▶  $\rho_\eta$  density matrix on  $\mathcal{H}_\eta$
- ▶  $\text{Tr}_{\mathcal{H}_{\eta' \rightarrow \eta}} \rho_{\eta'} = \rho_\eta$

[Projective state spaces: Kijowski '76, Okołów '09 & '13]

# Projective State Spaces

## Quantization

|                   | $\mathcal{M}_\eta \dashrightarrow \mathcal{H}_\eta$                                                           | $\mathcal{M}_{\eta'} \approx \mathcal{M}_{\eta' \rightarrow \eta} \times \mathcal{M}_\eta$<br>$\dashrightarrow \mathcal{H}_{\eta'} \approx \mathcal{H}_{\eta' \rightarrow \eta} \otimes \mathcal{H}_\eta$ |
|-------------------|---------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Holo.<br>repr.    | Holomorphic st. on $\mathcal{M}_\eta$<br>$\mathcal{H}_\eta = L_2(\mathcal{M}_\eta, dv_\eta) \cap \text{Holo}$ | $\exists$ holomorphic st.<br>on $\mathcal{M}_{\eta' \rightarrow \eta}$ such that<br>$\approx$ is holomorphic                                                                                              |
| Position<br>repr. | $\mathcal{M}_\eta = T^*(\mathcal{C}_\eta)$<br>$\mathcal{H}_\eta = L_2(\mathcal{C}_\eta, d\mu_\eta)$           | $\mathcal{C}_{\eta'} \approx \mathcal{C}_{\eta' \rightarrow \eta} \times \mathcal{C}_\eta$<br>$\mu_{\eta'} \approx \mu_{\eta' \rightarrow \eta} \times \mu_\eta$                                          |

+ observables...


[Geometric quantization: Woodhouse '92]

# Quantum Projective State Spaces

Example: Second Quantization

$$\mathcal{F}_{I'} \approx \mathcal{F}_{I^\perp \cap I'} \otimes \mathcal{F}_I$$

$$|n_1, n_2, n_3\rangle = |n_3\rangle \otimes |n_1, n_2\rangle$$



$$I \subset I' \subset \mathcal{H}_{\text{one particule}}$$

Projective system for the second quantization of  $\mathcal{H}_{\text{one particule}}$ :

- ▶ labels:  $I \subset \mathcal{H}_{\text{one particule}}$  with  $\dim I < \infty$
- ▶ 'small' Hilbert space:  
 $\mathcal{F}_I$  = partial Fock space built on  $I$
- ▶  $I \subset I' \Rightarrow \mathcal{F}_{I'} \approx \mathcal{F}_{I^\perp \cap I'} \otimes \mathcal{F}_I$

Mapping a density matrix  $\sigma$  on  $\mathcal{F}_{\text{Fock}}$  to a projective state  $(\rho_I)_I$ :

- ▶  $\forall I, \rho_I = \text{Tr}_{\mathcal{F}_{I^\perp}} \sigma$   
using  $\mathcal{F}_{\text{Fock}} \approx \mathcal{F}_{I^\perp} \otimes \mathcal{F}_I$

# Inductive Limit from a Choice of Vacuum Fock Space

Fock space  $\mathcal{F}_{\text{Fock}}$  built on  $\mathcal{H}_{\text{one particule}}$  as an inductive limit:

- ▶  $\emptyset_{I' \rightarrow I} = |(0)_j\rangle \in \mathcal{F}_{I^\perp \cap I'}$
- ▶  $\iota_{I' \leftarrow I} : |(n_i)_i\rangle \in \mathcal{F}_I \mapsto |(0)_j, (n_i)_i\rangle \in \mathcal{F}_{I'} \approx \mathcal{F}_{I^\perp \cap I'} \otimes \mathcal{F}_I$

Mapping a density matrix  $\sigma$  on  $\mathcal{F}_{\text{Fock}}$  to a projective state  $(\rho_I)_I$ :

- ▶  $\forall I \subset \mathcal{H}_{\text{one particule}}, \rho_I = \text{Tr}_{\mathcal{F}_{I^\perp}} \sigma$   
using the factorization  $\mathcal{F}_{\text{Fock}} \approx \mathcal{F}_{I^\perp} \otimes \mathcal{F}_I$
- ▶  $\sup_I \inf_{I' \supset I} \text{Tr}_{\mathcal{F}_{I'}} \left[ (|\emptyset_{I' \rightarrow I}\rangle \langle \emptyset_{I' \rightarrow I}| \otimes \mathbf{1}_{\mathcal{F}_I}) \rho_{I'} \right] = 1$

# Infinite Tensor Product

Infinite tensor product

$$\bigotimes_{f \in \Lambda} \mathcal{J}_f := \bigoplus_{[\Omega]_{\sim}} \mathcal{H}_{[\Omega]_{\sim}}$$

$$\Omega \simeq \Omega' \Leftrightarrow \sum_{f \in \Lambda} |\langle \Omega_f, \Omega'_f \rangle - 1| < \infty$$

Projective system:

- $\lambda \subset \Lambda$ ,  $\#\lambda < \infty$
- $\mathcal{H}_\lambda := \bigotimes_{f \in \lambda} \mathcal{J}_f$

ITP states  $\leftrightarrow$  projective states:

- each sector separately
- cross-sector correlations are lost
- not representable on the ITP:

$$\rho = \bigotimes_{f \in \Lambda} \left( \frac{|\Omega_f\rangle\langle\Omega_f| + |\Omega'_f\rangle\langle\Omega'_f|}{2} \right)$$

[ITP: von Neumann '39]

Projective State Spaces for QG (S. Lanéry)

└ Extra Slides


└ Other State Spaces

arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)

Cofinal Others

# Implementation of Gauge Constraints and Dynamics

## Nice Constraints



Restrictive requirements:


- ▶ orbits are projected on orbits  $\rightarrow \pi_\eta^{\text{DYN}}$  between reduced phase spaces
- ▶ compatible with symplect. structures

Dynamical projective system & transport maps:

- ▶ states to projective families of orbits
- ▶ observables

# Implementation of Gauge Constraints and Dynamics

## Nice Constraints



Restrictive requirements:


- ▶ orbits are projected on orbits  $\rightarrow \pi_\eta^{\text{DYN}}$  between reduced phase spaces
- ▶ compatible with symplect. structures

Dynamical projective system & transport maps:

- ▶ states to projective families of orbits
- ▶ observables

# Implementation of Gauge Constraints and Dynamics

## Nice Constraints



Restrictive requirements:


- ▶ orbits are projected on orbits  $\rightarrow \pi_\eta^{\text{DYN}}$  between reduced phase spaces
- ▶ compatible with symplect. structures

Dynamical projective system & transport maps:

- ▶ states to projective families of orbits
- ▶ observables

# Implementation of Gauge Constraints and Dynamics

## Nice Constraints



Restrictive requirements:


- ▶ orbits are projected on orbits  $\rightarrow \pi_\eta^{\text{DYN}}$  between reduced phase spaces
- ▶ compatible with symplect. structures

Dynamical projective system & transport maps:

- ▶ states to projective families of orbits
- ▶ observables

# Implementation of Gauge Constraints and Dynamics

## Nice Constraints



Restrictive requirements:


- ▶ orbits are projected on orbits  $\rightarrow \pi_\eta^{\text{DYN}}$  between reduced phase spaces
- ▶ compatible with symplect. structures

Dynamical projective system & transport maps:

- ▶ states to projective families of orbits
- ▶ observables

# Implementation of Gauge Constraints and Dynamics

## Nice Constraints



Restrictive requirements:


- ▶ orbits are projected on orbits  $\rightarrow \pi_{\eta}^{\text{DYN}}$  between reduced phase spaces
- ▶ compatible with symplect. structures

Dynamical projective system & transport maps:

- ▶ states to projective families of orbits
- ▶ observables

# Implementation of Gauge Constraints and Dynamics

## Nice Constraints



Restrictive requirements:


- ▶ orbits are projected on orbits  $\rightarrow \pi_\eta^{\text{DYN}}$  between reduced phase spaces
- ▶ compatible with symplect. structures

Dynamical projective system & transport maps:

- ▶ states to projective families of orbits
- ▶ observables

# Implementation of Gauge Constraints and Dynamics

## Unfitting Constraints



Successive approximations:


- ▶ labelled by  $\varepsilon \in \mathcal{E}$
- ▶ nice on smaller and smaller cofinal parts of  $\mathcal{L}$

Projections between approximated theories:

- ▶ dynamical projective system on a subset of  $\mathcal{E} \times \mathcal{L}$
- ▶ notion of convergence

# Implementation of Gauge Constraints and Dynamics

## Unfitting Constraints



Successive approximations:


- ▶ labelled by  $\varepsilon \in \mathcal{E}$
- ▶ nice on smaller and smaller cofinal parts of  $\mathcal{L}$

Projections between approximated theories:

- ▶ dynamical projective system on a subset of  $\mathcal{E} \times \mathcal{L}$
- ▶ notion of convergence

# Implementation of Gauge Constraints and Dynamics

## Unfitting Constraints



Successive approximations:


- ▶ labelled by  $\varepsilon \in \mathcal{E}$
- ▶ nice on smaller and smaller cofinal parts of  $\mathcal{L}$

Projections between approximated theories:

- ▶ dynamical projective system on a subset of  $\mathcal{E} \times \mathcal{L}$
- ▶ notion of convergence

# Implementation of Gauge Constraints and Dynamics

## Unfitting Constraints



Successive approximations:


- ▶ labelled by  $\varepsilon \in \mathcal{E}$
- ▶ nice on smaller and smaller cofinal parts of  $\mathcal{L}$

Projections between approximated theories:

- ▶ dynamical projective system on a subset of  $\mathcal{E} \times \mathcal{L}$
- ▶ notion of convergence

# Implementation of Gauge Constraints and Dynamics

## Unfitting Constraints



Successive approximations:

- ▶ labelled by  $\varepsilon \in \mathcal{E}$
- ▶ nice on smaller and smaller cofinal parts of  $\mathcal{L}$

Projections between approximated theories:

- ▶ dynamical projective system on a subset of  $\mathcal{E} \times \mathcal{L}$
- ▶ notion of convergence

# Implementation of Gauge Constraints and Dynamics

Toy Model: Schrödinger Equation

$\infty$

$$E - \langle \psi, H\psi \rangle = 0$$

$$\mathcal{M}_\infty^{\text{KIN}} = \mathcal{H}_{\text{one part.}} \times \mathbb{R}^2 = \{(\psi; t, E)\}$$

Approximations:

- ▶  $\epsilon > 0$  deformation → compact orbits
- ▶ truncation on finite dim. subspace  $I$

Proof of principle for previous strategy:

- ▶ classical → convergence for normed dynamical states
- ▶ quantum → convergence for Fock dynamical states

# Implementation of Gauge Constraints and Dynamics

Toy Model: Schrödinger Equation

$\infty$

$$E - \langle \psi, H\psi \rangle = 0$$

$$\mathcal{M}_\infty^{\text{KIN}} = \mathcal{H}_{\text{one part.}} \times \mathbb{R}^2 = \{(\psi; t, E)\}$$

$\epsilon > 0$

$$(E - \langle \psi, H\psi \rangle)^2 + \epsilon^4 t^2 = \epsilon^2$$

Approximations:

- ▶  $\epsilon > 0$  deformation → compact orbits
- ▶ truncation on finite dim. subspace  $I$

Proof of principle for previous strategy:

- ▶ classical → convergence for normed dynamical states
- ▶ quantum → convergence for Fock dynamical states

# Implementation of Gauge Constraints and Dynamics

Toy Model: Schrödinger Equation

$\infty$

$$E - \langle \psi, H\psi \rangle = 0$$

$$\mathcal{M}_\infty^{\text{KIN}} = \mathcal{H}_{\text{one part.}} \times \mathbb{R}^2 = \{(\psi; t, E)\}$$

$$\epsilon > 0, \quad I \subset \mathcal{H}_{\text{one part.}}$$

$$(E - \langle \psi, H_I \psi \rangle)^2 + \epsilon^4 t^2 = \epsilon^2$$

$$\& \quad \psi \in I$$

where  $H_I = \Pi_I H \Pi_I$

Approximations:



- ▶  $\epsilon > 0$  deformation  $\rightarrow$  compact orbits
- ▶ truncation on finite dim. subspace  $I$

Proof of principle for previous strategy:

- ▶ classical  $\rightarrow$  convergence for normed dynamical states
- ▶ quantum  $\rightarrow$  convergence for Fock dynamical states

# Implementation of Gauge Constraints and Dynamics

Toy Model: Schrödinger Equation



Approximations:



- ▶  $\epsilon > 0$  deformation  $\rightarrow$  compact orbits
- ▶ truncation on finite dim. subspace  $I$

Proof of principle for previous strategy:

- ▶ classical  $\rightarrow$  convergence for normed dynamical states
- ▶ quantum  $\rightarrow$  convergence for Fock dynamical states

# Implementation of Gauge Constraints and Dynamics

Toy Model: Schrödinger Equation



Approximations:

- ▶  $\epsilon > 0$  deformation  $\rightarrow$  compact orbits
- ▶ truncation on finite dim. subspace  $I$

Proof of principle for previous strategy:

- ▶ classical  $\rightarrow$  convergence for normed dynamical states
- ▶ quantum  $\rightarrow$  convergence for Fock dynamical states

# Quantizing Gravity \*

Perturbative  
Formulation

- ▶  $g = g_o + h$
- ▶ perturb. QFT

---

**non-**  
renormalizable

---

\* disclaimer: this table is **not** exhaustive!

[Non-renormalizability: Goroff & Sagnotti '85]

Projective State Spaces for QG (S. Lanéry)

└ Extra Slides

└ LQG Basics

arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)

# Quantizing Gravity \*

|                               |                      |  |
|-------------------------------|----------------------|--|
| Perturbative<br>Formulation   |                      |  |
| ► $g = g_o + h$               |                      |  |
| ► perturb. QFT                |                      |  |
| <b>non-</b><br>renormalizable |                      |  |
| String<br>Theory              | Asymptotic<br>Safety |  |

\* disclaimer: this table is **not** exhaustive!

[Strings: Green, Polyakov, Schwarz, Witten... '80s; Asympt. safe: Parisi, Weinberg, Wilson... '70s]

# Quantizing Gravity \*

|                               |                             |                                                 |
|-------------------------------|-----------------------------|-------------------------------------------------|
| Perturbative<br>Formulation   | Dynamical<br>Triangulations |                                                 |
| ► $g = g_o + h$               | ► discretization            |                                                 |
| ► perturb. QFT                | ► path-integral             |                                                 |
| <b>non-</b><br>renormalizable |                             | dominated by<br><b>degenerate</b><br>geometries |
| String<br>Theory              | Asymptotic<br>Safety        | Causal Dyn.<br>Triangl.                         |

\* disclaimer: this table is **not** exhaustive!

[(C)DT: Ambjørn, Jurkiewicz, Loll... '90s]

# Quantizing Gravity \*


|                                                                                                   |                                                                                          |                                                                                                 |
|---------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------|
| Perturbative Formulation                                                                          | Dynamical Triangulations                                                                 | Geometrodynamics                                                                                |
| <ul style="list-style-type: none"><li>▶ <math>g = g_o + h</math></li><li>▶ perturb. QFT</li></ul> | <ul style="list-style-type: none"><li>▶ discretization</li><li>▶ path-integral</li></ul> | <ul style="list-style-type: none"><li>▶ canonical GR</li><li>▶ canonical quantization</li></ul> |
| <b>non-renormalizable</b>                                                                         |                                                                                          | dominated by<br><b>degenerate geometries</b>                                                    |
| String Theory                                                                                     | Asymptotic Safety                                                                        | Causal Dyn. Triangl.                                                                            |

\* disclaimer: this table is **not** exhaustive!

[Geometrodynamics: DeWitt, Misner, Wheeler... '60s]

# Canonical General Relativity

## The ADM Formalism



[Canonical GR: Arnowitt, Deser, Misner '62]

Phase space of spacial slices:


- ▶ configuration variables:  
3-geometry
- ▶ momentum variables:  
extrinsic curvature

Gauge constraints:

- ▶ 3-slice in constraint surface  
iff can be sliced out of  
Einstein 4-geometry
- ▶ different 3-slices in same  
gauge orbit iff can be sliced  
out of the **same** Einstein  
4-geometry

# Canonical General Relativity

## The ADM Formalism



[Canonical GR: Arnowitt, Deser, Misner '62]

Phase space of spacial slices:


- ▶ configuration variables:  
3-geometry
- ▶ momentum variables:  
extrinsic curvature

Gauge constraints:

- ▶ 3-slice in constraint surface  
iff can be sliced out of  
Einstein 4-geometry
- ▶ different 3-slices in same  
gauge orbit iff can be sliced  
out of the **same** Einstein  
4-geometry

# Canonical General Relativity

## The ADM Formalism



[Canonical GR: Arnowitt, Deser, Misner '62]

Phase space of spacial slices:


- ▶ configuration variables:  
3-geometry
- ▶ momentum variables:  
extrinsic curvature

Gauge constraints:

- ▶ 3-slice in constraint surface  
iff can be sliced out of  
Einstein 4-geometry
- ▶ different 3-slices in same  
gauge orbit iff can be sliced  
out of the **same** Einstein  
4-geometry

# Canonical General Relativity

## The ADM Formalism



[Canonical GR: Arnowitt, Deser, Misner '62]

Phase space of spacial slices:

- ▶ configuration variables:  
3-geometry
- ▶ momentum variables:  
extrinsic curvature

Gauge constraints:

- ▶ 3-slice in constraint surface  
iff can be sliced out of  
Einstein 4-geometry
- ▶ different 3-slices in same  
gauge orbit iff can be sliced  
out of the **same** Einstein  
4-geometry

# Quantizing Gravity \*

|                                                                                                   |                                                                                          |                                                                                                 |
|---------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------|
| Perturbative Formulation                                                                          | Dynamical Triangulations                                                                 | Geometrodynamics                                                                                |
| <ul style="list-style-type: none"><li>▶ <math>g = g_o + h</math></li><li>▶ perturb. QFT</li></ul> | <ul style="list-style-type: none"><li>▶ discretization</li><li>▶ path-integral</li></ul> | <ul style="list-style-type: none"><li>▶ canonical GR</li><li>▶ canonical quantization</li></ul> |
| <b>non-renormalizable</b>                                                                         |                                                                                          | dominated by<br><b>degenerate geometries</b>                                                    |
| String Theory                                                                                     | Asymptotic Safety                                                                        | Causal Dyn. Triangl.                                                                            |

\* disclaimer: this table is **not** exhaustive!

[Geometrodynamics: DeWitt, Misner, Wheeler... '60s]

# Quantizing Gravity \*

|                           |                                           |                                     |
|---------------------------|-------------------------------------------|-------------------------------------|
| Perturbative Formulation  | Dynamical Triangulations                  | Geometrodynamics                    |
| ► $g = g_0 + h$           | ► discretization                          | ► canonical GR                      |
| ► perturb. QFT            | ► path-integral                           | ► canonical quantization            |
| <b>non-renormalizable</b> | <b>dominated by degenerate geometries</b> | <b>hard to explicitly implement</b> |
| String Theory             | Asymptotic Safety                         | Causal Dyn. Triangl.                |
|                           |                                           | Spin Foams                          |
|                           |                                           | Loop Quant. Gravity                 |

\* disclaimer: this table is **not** exhaustive!


[LQG/Spin foams: Ashtekar, Lewandowski, Pullin, Rovelli, Smolin, Thiemann... '80s]

# Loop Quantum Gravity


## The Holonomy-Flux Variables

$$E_S \in \text{Lie}^*(G)$$

$$h_e \in G$$



$$\{h_e, E_{S,i}\} = \tau_i \cdot h_e$$



$$\{E_{S,i}, E_{S',j}\} = C_{ij}^k E_{S \cap S', k}$$

GR as a  $G = SU(2)$  gauge theory:

- ▶  $\text{Lie}^*(G)$ -valued **fluxes** encode spacial 3-geometry
- ▶  $G$ -valued **holonomies** combine intrinsic and extrinsic curvature
- ▶ Poisson algebra can be regularized, no background metric needed
- ▶ reduces to ADM formalism once additional  $G$ -gauge invariance is imposed

[Holonomy-flux algebra: Ashtekar, Isham, Rovelli, Smolin, Lewandowski, Pullin, Gambini,...]


# Loop Quantum Gravity

## The Ashtekar–Lewandowski Hilbert Space

### Fock Space

$$I = \text{span} \{ e_1, e_2 \}$$

$$\subset \mathcal{H}_{\text{one particle}}$$



$$\mathcal{F}_I = L_2(\mathbb{R}) \otimes L_2(\mathbb{R})$$

$$= \text{span} \{ |n_1, n_2\rangle \mid n_1, n_2 \in \mathbb{N} \}$$

$$\hat{a}_{e_1} |n_1, n_2\rangle = \sqrt{n_1} |n_1 - 1, n_2\rangle$$


$$\hat{a}_{e_1}^\dagger |n_1, n_2\rangle = \sqrt{n_1 + 1} |n_1 + 1, n_2\rangle$$

[LQG Hilbert space: Isham, Ashtekar, Lewandowski,... '92–95]

# Loop Quantum Gravity


The Ashtekar–Lewandowski Hilbert Space

Fock Space



$$\begin{aligned} I &= \text{span } \{e_1, e_2\} \\ &\subset \mathcal{H}_{\text{one particle}} \end{aligned}$$

AL Hilbert Space




$$\begin{aligned} \gamma &= (e_1, e_2) \\ &\text{on the spacial slice } \Sigma \end{aligned}$$

$$\begin{aligned} \mathcal{F}_I &= L_2(\mathbb{R}) \otimes L_2(\mathbb{R}) \\ &= \text{span} \{ |n_1, n_2\rangle \mid n_1, n_2 \in \mathbb{N}\} \end{aligned}$$

$$\begin{aligned} \mathcal{H}_\gamma &= L_2(G) \otimes L_2(G) \\ &= \{ |\psi_\gamma\rangle \mid \psi_\gamma : g_1, g_2 \mapsto \psi_\gamma(g_1, g_2)\} \end{aligned}$$

$$\begin{aligned} \hat{a}_{e_1} |n_1, n_2\rangle &= \sqrt{n_1} |n_1 - 1, n_2\rangle \\ \hat{a}_{e_1}^\dagger |n_1, n_2\rangle &= \sqrt{n_1 + 1} |n_1 + 1, n_2\rangle \end{aligned}$$

$$\begin{aligned} \widehat{f \circ h}_{e_1} |\psi_\gamma\rangle &= |f(g_1) \psi_\gamma\rangle \\ \widehat{E}_{S,i} |\psi_\gamma\rangle &= |i \nabla_{\tau_i}^{[g_1]} \psi_\gamma\rangle \end{aligned}$$



[LQG Hilbert space: Isham, Ashtekar, Lewandowski,... '92–95]

Projective State Spaces for QG (S. Lanéry)

└ Extra Slides

└ LQG Basics


arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)

Others InjAL

# Loop Quantum Gravity

## The Ashtekar–Lewandowski Hilbert Space

### Fock Space




$$I' = \text{span} \{e_1, e_2, e_3\}$$

$$I = \text{span} \{e_1, e_2\}$$

$\subset \mathcal{H}_{\text{one particle}}$

### AL Hilbert Space



$$\gamma = (e_1, e_2)$$

on the spacial slice  $\Sigma$

$$\mathcal{F}_I = \text{span} \{ |n_1, n_2\rangle \mid n_1, n_2 \in \mathbb{N}\}$$

$$\mathcal{H}_\gamma = \{ |\psi_\gamma\rangle \mid \psi_\gamma : g_1, g_2 \mapsto \psi_\gamma(g_1, g_2)\}$$

$$\mathcal{F}_I \subset \mathcal{F}_{I'}$$

$$|n_1, n_2\rangle \simeq |n_1, n_2, 0\rangle$$

$$\mathcal{F}_{\text{Fock}} = \lim_{I \subset \mathcal{H}_{\text{one part.}}} \mathcal{F}_I$$

[LQG Hilbert space: Isham, Ashtekar, Lewandowski,... '92–95]

Projective State Spaces for QG (S. Lanéry)

└ Extra Slides

└ LQG Basics


arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)

Others InjAL

# Loop Quantum Gravity

## The Ashtekar–Lewandowski Hilbert Space

### Fock Space




$$I' = \text{span} \{e_1, e_2, e_3\}$$

$$I = \text{span} \{e_1, e_2\}$$

$\subset \mathcal{H}_{\text{one particle}}$

### AL Hilbert Space



$$\gamma' = (e_1, e_2, e_3)$$

$$\gamma = (e_1, e_2)$$

on the spacial slice  $\Sigma$

$$\mathcal{F}_I = \text{span} \{ |n_1, n_2\rangle \mid n_1, n_2 \in \mathbb{N}\}$$

$$\mathcal{H}_\gamma = \{ |\psi_\gamma\rangle \mid \psi_\gamma : g_1, g_2 \mapsto \psi_\gamma(g_1, g_2)\}$$

$$\mathcal{F}_I \subset \mathcal{F}_{I'}$$

$$|n_1, n_2\rangle \simeq |n_1, n_2, 0\rangle$$

$$\mathcal{F}_{\text{Fock}} = \lim_{I \subset \mathcal{H}_{\text{one part.}}} \mathcal{F}_I$$

$$\mathcal{H}_\gamma \subset \mathcal{H}_{\gamma'}$$


$$|\psi_\gamma\rangle \simeq |\psi_{\gamma'} : g_1, g_2, g_3 \mapsto \psi_\gamma(g_1, g_2)\rangle$$

$$\mathcal{H}_{\text{AL}} = \lim_{\gamma \text{ on } \Sigma} \mathcal{H}_\gamma$$

[LQG Hilbert space: Isham, Ashtekar, Lewandowski,... '92–95]

# Projective State Space for LQG

## The Label Set




The label set:

- ▶ a graph = a choice of holonomies
- ▶ a set of surfaces dual to this graph = a choice of conjugate fluxes
- ▶ the label set must be directed  
⇒ allow intersecting surfaces
- ▶ each label = a  $\{\cdot, \cdot\}$ -subalgebra  
⇒ need edges probing any intersection

[See also: Okołów '13 (Abelian case)]

# Projective State Space for LQG

## The Label Set




The label set:

- ▶ a graph = a choice of holonomies
- ▶ a set of surfaces dual to this graph = a choice of conjugate fluxes
- ▶ the label set must be directed  
⇒ allow intersecting surfaces
- ▶ each label = a  $\{\cdot, \cdot\}$ -subalgebra  
⇒ need edges probing any intersection

[See also: Okołów '13 (Abelian case)]

# Projective State Space for LQG

## The Label Set




The label set:

- ▶ a graph = a choice of holonomies
- ▶ a set of surfaces dual to this graph = a choice of conjugate fluxes
- ▶ the label set must be directed  
⇒ allow intersecting surfaces
- ▶ each label = a  $\{\cdot, \cdot\}$ -subalgebra  
⇒ need edges probing any intersection

[See also: Okołów '13 (Abelian case)]


# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )





# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )





# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )






# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )






# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )







# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )







# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )








# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )








# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )




# Holonomy-Flux Algebra

Directedness of the Label Set ( $\forall \eta, \eta' \in \mathcal{L}, \exists \eta'' \succcurlyeq \eta, \eta'$ )



# Universal Sequences: A 1-dimensional Model

## Selection of an Increasing Sequence



Simplified version of holonomy-flux algebra:

- ▶ one-dimensional
- ▶ one-sided fluxes, acting only on the left


Selecting an increasing  $(\eta_k)_{k \in \mathbb{N}}$ :

- ▶  $\eta_k = \{e_{\ell/2^k}, S_{\ell/2^k} \mid 1 \leq \ell \leq 2^k\}$
- ▶  $k \leq k' \Rightarrow \eta_k \preceq \eta_{k'}$

[See also: combinatorial LQG: Zapata '97; discrete quantum gravity: Gambini & Pullin;...]

# Universal Sequences: A 1-dimensional Model

## Universality and Approximation of Diffeomorphisms



One can formulate generic properties that  $(\eta_k)_k$  should satisfy.


All sequences with these properties are equivalent:

- ▶ deformation mapping  
eg.  $\{n/3^m\}$  to  $\{\ell/2^k\}$ , arbitrarily close to identity
- ▶ ⇒ identification of the corresponding state spaces
- ▶ similarly: approximating diffeomorphisms

[≠ universality at diffeo.-inv. level: Baez & Sawin '95, Zapata '97,...]

# Universal Sequences: A 1-dimensional Model

## Universality and Approximation of Diffeomorphisms



One can formulate generic properties that  $(\eta_k)_k$  should satisfy.


All sequences with these properties are equivalent:

- ▶ deformation mapping  
eg.  $\{n/3^m\}$  to  $\{\ell/2^k\}$ , arbitrarily close to identity
- ▶ ⇒ identification of the corresponding state spaces
- ▶ similarly: approximating diffeomorphisms

[≠ universality at diffeo.-inv. level: Baez & Sawin '95, Zapata '97,...]

# Universal Sequences: A 1-dimensional Model

## Universality and Approximation of Diffeomorphisms



One can formulate generic properties that  $(\eta_k)_k$  should satisfy.


All sequences with these properties are equivalent:

- ▶ deformation mapping  
eg.  $\{n/3^m\}$  to  $\{\ell/2^k\}$ , arbitrarily close to identity
- ▶ ⇒ identification of the corresponding state spaces
- ▶ similarly: approximating diffeomorphisms

[≠ universality at diffeo.-inv. level: Baez & Sawin '95, Zapata '97,...]

# Universal Sequences: A 1-dimensional Model

## Universality and Approximation of Diffeomorphisms



One can formulate generic properties that  $(\eta_k)_k$  should satisfy.


All sequences with these properties are equivalent:

- ▶ deformation mapping  
eg.  $\{n/3^m\}$  to  $\{\ell/2^k\}$ , arbitrarily close to identity
- ▶ ⇒ identification of the corresponding state spaces
- ▶ similarly: approximating diffeomorphisms

[≠ universality at diffeo.-inv. level: Baez & Sawin '95, Zapata '97,...]

# Universal Sequences: A 1-dimensional Model

## Universality and Approximation of Diffeomorphisms



One can formulate generic properties that  $(\eta_k)_k$  should satisfy.


All sequences with these properties are equivalent:

- ▶ deformation mapping  
eg.  $\{n/3^m\}$  to  $\{\ell/2^k\}$ , arbitrarily close to identity
- ▶ ⇒ identification of the corresponding state spaces
- ▶ similarly: approximating diffeomorphisms

[≠ universality at diffeo.-inv. level: Baez & Sawin '95, Zapata '97,...]

# Obstruction to Finite Variances in the $G = \mathbb{R}$ Case

Selecting a Set of Observables with Uniformly Bounded Variances



Looking for a state such that:


- $\forall e, \Delta h_e^2 < \infty$
- $\forall S, \Delta E_S^2 < \infty$

$\{p \in e\}$  uncountable:

- $\exists A /$   
 $\{p \mid \Delta h_{e_p}^2 \leq A\}$  uncountable
- $\exists A, B /$   
 $\{p \mid \Delta h_{e_p}^2 \leq A, \Delta E_{S_p}^2 \leq B\}$  uncountable

# Obstruction to Finite Variances in the $G = \mathbb{R}$ Case

Bound from Heisenberg Uncertainty Relations



One can construct a quadratic operator  $Q_N$  from:

- ▶  $h_{e_1}, \dots, h_{e_N}$
- ▶  $E_{S_1}, \dots, E_{S_N}$

such that  $\langle Q_N \rangle \leq A + B$

But Heisenberg inequalities  
⇒ **lower** bound on  $\langle Q_N \rangle$  that  
**diverges** with  $N!$

[See also: Varadarajan; Lewandowski, Okołów, Sahlmann, Thiemann; Fleischhacker;...]

**Introduction**  
Quantum Field Theory without a Vacuum State

**Standard approach to QFT:**

- discrete excitations around a **vacuum state**
- if **vacuum state is unstable** -> **inconsistency** -> need to find the **state to the dynamics**
- QFT on curved space time is **not natural choice**
- **LQG**: performed **background subtraction** (vacuum state)
- **loop quantum gravity**: **quantum state** and the **dynamics**, and the semi-classical limit (see [Ashtekar, Bojowald & Schenkel](#))

An alternative way to construct the state space ([Pfenning '96, Grotz '04](#))

- by going along highly elementary building blocks
- **spin networks** (labeled by **bundles** shown to QFT)
- **global universal quantum state**, independent of any choice of polarization

**Projective Formalism for QFT**  
Contracting between Fock Spaces

$M_A = M_{A_1} \otimes \dots \otimes M_{A_n}$



$M_B = M_{B_1} \otimes \dots \otimes M_{B_n}$

Collection of partial tensors:

- $\eta = \eta^L$  is a selection of  $d_A$ 's
- "real" space partial  $M_\eta$

Classical gauges:

- projections  $\eta \rightarrow \eta'$  for  $\eta \in \mathcal{L}$
- linking observables
- $A_\eta = A_\eta^L$  compatible with the Poisson brackets
- general factorization
- $M_\eta = M_{\eta^L} \otimes M_{\eta^R}$

Projection into space: Kijowski, Henneaux, 90 & 10f

Diagram illustrating the tensor product structure of  $M_A$  and  $M_B$ , showing a large square divided into a grid of smaller squares.

The diagram illustrates the Projective Formalism for Quantum Field Theory (QFT). It shows the relationship between the Projective State Space ( $\mathcal{S}_P$ ) and the Hilbert Space ( $\mathcal{H}$ ). The Projective State Space is defined as the set of equivalence classes of states  $[|\psi\rangle]$ , where two states are equivalent if their difference is in the kernel of the energy operator ( $\mathcal{K}_{E_0}$ ). This leads to a quotient space structure. The Hilbert Space  $\mathcal{H}$  is shown as a subspace of the Projective State Space. A map  $\pi$  from the Projective State Space to the Hilbert Space is defined by  $\pi([\psi]) = |\psi\rangle$ . The image of this map is denoted as  $\mathcal{H}_P$ . The Hilbert Space  $\mathcal{H}$  is also shown as a subspace of the Projective State Space. A map  $\pi_P$  from the Projective State Space to the Hilbert Space is defined by  $\pi_P([\psi]) = |\psi\rangle$ . The image of this map is denoted as  $\mathcal{H}_H$ .

**Universal Selections of Degrees of Freedom**

Reservoir as a Quasi-Critical System of Lattes

Continuous de la C.V.

- abstraction for math convenience
- direct countable sub-objects of algebras is enough — basic
- universal selection and respect properties of the theory

Quasi-critical increasing temperature:

Up to  $L_c = 3n/2$

- $a \leq n \leq m$  (with  $m = m_0$ )
- the (any  $m \geq 1$ )
- the (any  $m \geq 1$ )

**Universal Selections of Degrees of Freedom**

Universality & Symmetries

Subsequences = deformations /

- $\{u_{1,1}\} \subset \{u_{1,n}\}$
- $\{u_{1,1}\} \subset \{u_{2,n}\}$
- $\{u_{1,1}\} \subset \{u_{3,n}\}$
- $\{u_{1,1}\} \subset \{u_{4,n}\} \cup \{u_{5,n}\}$
- $\{u_{1,1}\} \subset \{u_{4,n}\} \cup \{u_{5,n}\}$

Approximating symmetries of deformation

- $\{u_{1,1}\} \subset \{u_{1,n}\}$
- $\{u_{1,1}\} \subset \{u_{2,n}\}$
- $\{u_{1,1}\} \subset \{u_{3,n}\}$
- $\{u_{1,1}\} \subset \{u_{4,n}\} \cup \{u_{5,n}\}$
- (with  $\Phi = e^{-\lambda x^2}$  and  $\Psi = e^{-\lambda y^2}$ )

| Summary:                 | Comparison with other State Spaces |                      |                               |       |             |     | ITP |
|--------------------------|------------------------------------|----------------------|-------------------------------|-------|-------------|-----|-----|
|                          | nodes<br>holes & flows             | refined<br>grad. AOG | refined<br>grad. sub-algebras | index | proj.<br>on | ITP |     |
| Hilbert space            | ✓                                  | ✓                    | ✓                             | ✓     | ✓           | ✓   | ✓   |
| Separability             | ✓                                  | ✗                    | ✓                             | ✓     | ✓           | ✓   | ✓   |
| Universality             | ✓                                  | ✓                    | ✓                             | ✓     | ✓           | ✓   | ✓   |
| Axioms of choice         | ✓                                  | ✓                    | ✓                             | ✓     | ✓           | ✓   | ✓   |
| Algebraic semantics      | ✓                                  | ✓                    | ✓                             | ✓     | ✓           | ✓   | ✓   |
| No-compact G             | ✗                                  | ✗                    | ✗                             | ✗     | ✗           | ✗   | ✗   |
| Universal quantification | ✓                                  | ✓                    | ✓                             | ✓     | ✓           | ✓   | ✓   |

[ITP] von Neumann AOG, Gödel & Thurston [6]. [2]

**Quantum Projective State Spaces**

Example:  $\text{H}_2$  molecule

$P = P_{\text{H}_2} \otimes P_{\text{H}_2}$

$[m_1, m_2] = [m_1] \otimes [m_2]$

$P \in \mathbb{C}^4 \otimes \mathbb{C}^4 \subset \mathbb{C}^{16}$

$\mathbb{C}^{16} \subset \mathbb{C}^{N \times N}$  (Nonsingular)

Projective system for the second quantization of  $N_m$  particles

- Labels:  $C_i$ :  $N_m$  particles
- States:  $|C_i\rangle$
- Variables:  $E_{ij}$ : partial Fock space label on  $i$
- Operators:  $T_{ij} = T_{i1} \otimes T_{j2} \otimes \dots \otimes T_{iN_m}$
- Mapping: a density matrix  $\rho$  on  $\mathcal{F}_{\text{H}_2}$  is a projective state  $\{\rho_C\}$
- Properties:  $\forall i, j: T_{ij} \rho_C T_{ij}^\dagger = \rho_C$
- along  $\sum_C \rho_C = I$

Implementation of Gauge Constraints and Dynamics

Reductive requirements

- orbits are projected on
- reduced phase space
- compatible with symplectic structure

Gauge constraints & transport maps

- states to project
- families of orbits
- observables

The figure consists of two parts. The top part shows a coordinate system with axes x and y, and a red curve representing a trajectory. A vector field is shown as a series of arrows along the curve. The bottom part shows a similar coordinate system with axes x and y, but the curve is now red and wavy, representing a deformed trajectory. A vector field is also shown along this curve.

Projective State Spaces for QG (S. Lanéry)

## └ Extra Slides

L No-Go

arXiv: 1604.05629 & 1411.3592 (with T. Thiemann)